

87-265660 B4 D16

8



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

(11) Publication number:

0 238 323  
A2

(12)

## EUROPEAN PATENT APPLICATION

(21) Application number: 87302318.8

(22) Date of filing: 18.03.87

(51) Int. Cl.<sup>3</sup>: C 12 N 15/00  
C 12 N 9/00, C 12 N 1/20  
C 12 P 19/62  
/(C12N1/20, C12R1:19, 1:465)

INCREASING ANTIBIOTIC PROD. IN MICROORGANISMS +  
BY TRANSFORMING WITH A DNA CLONING VECTOR CODING  
FOR EXPRESSION OF A RATE-LIMITING ENZYME OR PROD. IN B

(30) Priority: 21.03.86 US 842330  
25.07.86 US 890670

(43) Date of publication of application:  
23.09.87 Bulletin 87/39

(54) Designated Contracting States:  
AT BE CH DE ES FR GB GR IT LI LU NL SE

(71) Applicant: ELI LILLY AND COMPANY  
Lilly Corporate Center  
Indianapolis Indiana 46285(US)

(72) Inventor: Cox, Karen Leigh  
Box 204 Painted Hills  
Martinsville Indiana 46151(US)

(72) Inventor: Fishman, Scott Eric  
5871 Hillside Avenue  
Indianapolis Indiana 46220(US)

(72) Inventor: Harshbarger, Charles Lee  
Rural Route No. 1, Box 343  
New Palestine Indiana 46163(US)

(72) Inventor: Seno, Eugene Thomas  
5109 East 72nd Street  
Indianapolis Indiana 46250(US)

(74) Representative: Hudson, Christopher Mark et al,  
Erl Wood Manor  
Windlesham Surrey GU20 6PH(GB)

(54) Improvements in or relating to antibiotic-producing microorganisms.

(57) A means for increasing the antibiotic-producing ability of an antibiotic-producing microbial host cell is disclosed. The method involves transforming an antibiotic-producing microorganism with a recombinant DNA cloning vector that codes for the expression of a rate-limiting antibiotic biosynthetic enzyme or other gene product.

EP 0 238 323 A2

2 2 1 9

87265660

Improvements In or Relating To  
Antibiotic-Producing Microorganisms

5 The present invention provides a novel method  
for increasing the antibiotic-producing ability of an  
antibiotic-producing organism. The method involves  
transforming a microbial host cell with a DNA sequence  
that codes for the expression of a gene product that is  
rate-limiting in the desired antibiotic's biosynthetic  
10 pathway. The invention also provides related DNA  
sequences that code for antibiotic biosynthetic gene  
products, recombinant DNA expression vectors, and  
transformed microbial host cells.

15 The present invention represents an early and  
significant commercial exploitation of recombinant DNA  
technology in antibiotic-producing organisms such as  
streptomycetes. Prior to the present invention, the  
development and exploitation of recombinant DNA technol-  
ogy has been limited, for the most part, to the expres-  
20 sion of specific polypeptides in E. coli and, in some  
instances, mammalian cells. These advances led to the  
comparatively simple expression of heterologous gene  
products such as human insulin A and B chains, human  
proinsulin, human growth hormone, human protein C, human  
25 tissue plasminogen activator, bovine growth hormone, and  
several other compounds of potential value. In each  
case, heterologous gene expression is more or less  
independent and does not interact with, take part in, or  
modulate operative biosynthetic pathways. Recombinant  
30 DNA technology can now be applied to improve selected

biosynthetic pathways for the expression of increased yields of antibiotics or antimicrobial precursors.

Most recombinant DNA technology applied to streptomycetes and other antibiotic-producing organisms has been limited to the development of cloning vectors. Early attempts include the disclosures of Reusser U.S. Patent No. 4,332,898 and Manis et al. U.S. Patent Nos. 4,273,875; 4,332,900; 4,338,400; and 4,340,674. Transformation of streptomycetes was not disclosed or taught in these early references. Improved vectors showing greater potential for use in antibiotic-producing organisms were disclosed, for example, by Fayerman et al. in U.S. Patent No. 4,513,086; and Nakatsukasa et al. in U.S. Patent Nos. 4,513,085 and 4,416,994. These improved vectors contain markers that are selectable in streptomycetes, can be used to transform many important Streptomyces strains, and constitute the tools required for conducting more complicated gene cloning experiments.

One such experiment was recently reported by Hopwood et al., 1985, in Nature 314:642. Although Hopwood et al. reported the production of novel hybrid antibiotic pigments, the disclosure does not focus on increasing the antibiotic-producing ability or biosynthetic efficiency of a given host cell but instead describes the transferring of actinorhodin pigment biosynthetic genes from one Streptomyces strain to another.

The present invention is particularly useful because it allows commercial application of recombinant

DNA technology to streptomycetes and other antibiotic-producing organisms. Because over half of the clinically important antibiotics are produced by streptomycetes, it is especially desirable to develop methods that are applicable to these organisms. The present invention provides such methods and allows for the cloning of genes both for increasing the antibiotic-producing ability as well as for the production of new antibiotics and antibiotic precursors in an antibiotic-producing organism.

For purposes of the present invention, the following terms are as defined:

Antibiotic - a substance produced by a microorganism that, either naturally or with limited chemical modification, inhibits or prevents the growth of or kills another microorganism or eukaryotic cell.

Antibiotic Biosynthetic Gene - a DNA segment that encodes an enzymatic activity or encodes a product that regulates expression of an enzymatic activity and which is necessary for an enzymatic reaction for converting primary metabolites to antibiotic intermediates, which also may possess antibiotic activity, and perhaps then to antibiotics.

Antibiotic Biosynthetic Pathway - the entire set of antibiotic biosynthetic genes and biochemical reactions necessary for the process of converting primary metabo-

lites to antibiotic intermediates and then to antibiotics.

5 Antibiotic-Producing Microorganism - any organism, including, but not limited to Actinoplanes, Actinomadura, Bacillus, Cephalosporium, Micromonospora, Penicillium, Nocardia, and Streptomyces, that either produces an antibiotic or contains genes that, if expressed, would produce an antibiotic.

10 Antibiotic Resistance-Confering Gene - a DNA segment that encodes an activity that confers resistance to an antibiotic.

15 ApR - the ampicillin-resistant phenotype or gene conferring same.

20 Host Cell - an organism, including a viable protoplast thereof, which can be transformed with a recombinant DNA cloning vector.

NmR - the neomycin-resistant phenotype or gene conferring same.

25 Operation of Antibiotic Biosynthetic Pathway - the expression of antibiotic biosynthetic genes and the related biochemical reactions required for the conversion of primary metabolites into antibiotics.

Recombinant DNA Cloning Vector - any selectable and autonomously replicating or chromosomally integrating agent, including but not limited to plasmids and phages, comprising a DNA molecule to which additional DNA can be  
5 or has been added.

rep - as used in the Figures, a Streptomyces plasmid origin of replication.

10 Restriction Fragment - any linear DNA generated by the action of one or more restriction enzymes.

Sensitive Host Cell - a host cell, including a viable protoplast thereof, which cannot grow in the presence of  
15 a given antibiotic without a DNA segment that confers resistance thereto.

Transformant - a recipient host cell, including a viable protoplast thereof, which has undergone transformation.

20 Transformation - the introduction of DNA into a recipient host cell, including a viable protoplast thereof, that changes the genotype of the recipient cell.

25 tsr - the thiostrepton-resistant phenotype or gene conferring same.

The plasmid and chromosomal maps depicted in the Figures are drawn approximately to scale. However,  
30 the tylosin biosynthetic genes, although linked, are

scattered across a large segment of DNA. Therefore, detailed restriction site mapping data exists only for small regions of the large tylosin biosynthetic gene-containing DNA fragment. The maps do not necessarily provide an exhaustive listing of all the cut sites of a given restriction enzyme. The location of individual genes, represented by line segments on the maps, was determined by deletion mapping and thus only approximates the exact location of a given gene.

10           Figure 1 - The Tylosin Biosynthetic Pathway.

            Figure 2 - Restriction Site and Function Map of Plasmid pHJL280.

            Figure 3 - Restriction Site and Function Map of Plasmid pHJL284.

15           Figure 4 - Restriction Site and Function Map of Plasmid pHJL309.

            Figure 5 - Restriction Site and Function Map of Plasmid pHJL311.

20           Figure 6 - Restriction Site and Function Map of Plasmid pHJL315.

            Figure 7 - Chromosomal Organization of the Tylosin Biosynthetic Genes.

25           The present invention provides a method for increasing the antibiotic- or antibiotic precursor-producing ability of an antibiotic-producing microorganism, which comprises culturing a microorganism which produces an antibiotic or antibiotic precursor by a biosynthetic pathway, said microorganism being transformed with a DNA  
30           cloning vector or portion thereof which contains an

- antibiotic or antibiotic-precursor biosynthetic gene coding for expression of a rate-limiting enzyme or gene product of the biosynthetic pathway, under conditions suitable for cell growth, expression of the antibiotic
- 5 or antibiotic-precursor biosynthetic gene and production of the antibiotic or antibiotic precursor, provided that the culturing process provides an increase in the antibiotic-producing ability of the microorganism.
- 10 The invention further provides related antibiotic biosynthetic genes, recombinant DNA cloning vectors, and antibiotic or antibiotic precursor-producing microorganisms transformed with the genes and vectors.
- 15 Further, there is provided a process for preparing an antibiotic, an antibiotic precursor, or a pharmaceutically acceptable salt thereof, which comprises culturing a microorganism which produces an antibiotic or antibiotic precursor through an antibiotic biosynthetic
- 20 pathway, said microorganism being transformed with a DNA cloning vector, or portion thereof, in a culture medium containing assimilable sources of carbon, nitrogen and inorganic salts under aerobic fermentation conditions characterized in that the DNA cloning vector, or portion
- 25 thereof, comprises an antibiotic biosynthetic gene which codes for the expression of a rate-limiting enzyme or gene product of the antibiotic biosynthetic pathway, said antibiotic biosynthetic gene being expressed under fermentation conditions providing for an increase in the
- 30 antibiotic-producing ability of the microorganism.



The methods of the present invention are widely applicable to all antibiotic-producing organisms. The following tables provide a non-exhaustive list of antibiotic producing organisms to which the present invention may

5 apply.

X-6808A

-9-

TABLE I

## Aminocyclitol Antibiotic-Producing Organisms

| 5  | <u>Organism</u>                      | <u>Antibiotic</u>         |
|----|--------------------------------------|---------------------------|
|    | <u>Bacillus</u>                      |                           |
|    | various species                      | various aminocyclitols    |
| 10 | <u>Micromonospora</u>                |                           |
|    | various species                      | gentamycins               |
|    | <u>Saccharopolyspora</u>             |                           |
|    | various species                      | various aminocyclitols    |
| 15 | <u>Streptomyces</u>                  |                           |
|    | <u>albogriseolus</u>                 | neomycins                 |
|    | <u>albus</u> var. <u>metamycinus</u> | metamycin                 |
|    | <u>aquacanus</u>                     | N-methyl hygromycin B     |
| 20 | <u>atrofaciens</u>                   | hygromycins               |
|    | <u>bikiniensis</u>                   | streptomycin              |
|    | <u>bluensis</u> var. <u>bluensis</u> | bluensomycin              |
|    | <u>canus</u>                         | ribosyl paromamine        |
|    | <u>catenulae</u>                     | catenulin                 |
| 25 | <u>chrestomyceticus</u>              | aminosidine               |
|    | <u>crystallinus</u>                  | hygromycin A              |
|    | <u>erythrochromogenes</u>            |                           |
|    | var. <u>narutoensis</u>              | streptomycin              |
|    | <u>eurocidicus</u>                   | Al6316-C                  |
| 30 | <u>fradiae</u>                       | hybrimycins and neomycins |
|    | <u>fradiae</u> var. <u>italicus</u>  | aminosidine               |

X-6808A

-10-

TABLE I (Continued)

|    | <u>Organism</u>            | <u>Antibiotic</u>                               |
|----|----------------------------|---|
| 5  | <u>Streptomyces</u>        |   |
|    | <u>galbus</u>              | streptomycin                                    |
|    | <u>griseus</u>             | streptomycin                                    |
|    | <u>griseoflavus</u>        | MA 1267   |
|    | <u>hofuensis</u>           | seldomycin complex                              |
| 10 | <u>hygroscopicus</u>       | hygromycins,<br>leucanicidin, and<br>hygrolidin |
|    | <u>hygroscopicus</u> forma |   |
|    | <u>glebosus</u>            | glebomycin                                      |
| 15 | <u>hygroscopicus</u> var.  |   |
|    | <u>limoneus</u>            | validamycins                                    |
|    | <u>hygroscopicus</u> var.  |   |
|    | <u>sagamiensis</u>         | spectinomycin                                   |
|    | <u>kanamyceticus</u>       | kanamycin A and B                               |
| 20 | <u>kasugaensis</u>         | kasugamycins                                    |
|    | <u>kasugaspinus</u>        | kasugamycins                                    |
|    | <u>lavendulae</u>          | neomycin  |
|    | <u>lividus</u>             | lividomycins                                    |
|    | <u>mashuensis</u>          | streptomycin                                    |
| 25 | <u>microsporeus</u>        | SF-767  |
|    | <u>netropsis</u>           | LL-AM31   |
|    | <u>noboritoensis</u>       | hygromycins                                     |
|    | <u>olivaceus</u>           | streptomycin                                    |
|    | <u>olivoreticuli</u> var.  |   |
| 30 | <u>cellulophilus</u>       | destomycin A                                    |

X-6808A

-11-

TABLE I (Continued)

|    | <u>Organism</u>          | <u>Antibiotic</u>             |
|----|--------------------------|-------------------------------|
| 5  | <u>poolensis</u>         | streptomycin                  |
|    | <u>rameus</u>            | streptomycin                  |
|    | <u>ribosidificus</u>     | SF733                         |
|    | <u>rimofaciens</u>       | destomycin A                  |
|    | <u>rimosus</u> forma     |                               |
| 10 | <u>paromomycinus</u>     | paromomycins and<br>catenulin |
|    | <u>spectabilis</u>       | spectinomycin                 |
|    | <u>tenebrarius</u>       | tobramycin and<br>spramycin   |
| 15 | <u>Streptovercillium</u> |                               |
|    | <u>flavopersicus</u>     | spectinomycin                 |

X-6808A

-12-

TABLE II

## Ansamycin Antibiotic-Producing Organisms

| 5  | <u>Organism</u>                           | <u>Antibiotic</u>   |
|----|---|---|
|    | <u>Micromonospora</u>                     |   |
|    | various species                           | various ansamycins  |
| 10 | <u>Nocardia</u>                           |   |
|    | <u>mediterranei</u>                       | rifamycin   |
|    | <u>Streptomyces</u>                       |   |
|    | <u>collinus</u>                           | ansatrienes and<br>naphthomycins                                |
| 15 | <u>diastochromogenes</u>                  | ansatrienes and<br>naphthomycins                                |
|    | <u>galbus</u> subsp. <u>griseosporus</u>  | naphthomycin B  |
|    | <u>hygroscopicus</u>                      | herbimycin  |
| 20 | <u>hygroscopicus</u> var. <u>geldanus</u> |   |
|    | var. <u>nova</u>                          | geldamycin  |
|    | <u>nigellus</u>                           | 21-hydroxy-25-demethyl<br>25-methylthioproteo<br>streptovaricin |
| 25 | <u>rishiriensis</u>                       | mycotrienes   |
|    | sp. E/784                                 | actamycin and mycotrienes                                       |
|    | sp. E88                                   | mycotrienes   |
|    | <u>spectabilis</u>                        | streptovaricins   |
|    | <u>tolypophorous</u>                      | tolypomycin   |
| 30 |   |   |

TABLE III

## Anthracycline and Quinone Antibiotic-Producing Organisms

|    | <u>Organism</u>          | <u>Antibiotic</u>                                   |
|----|--------------------------|---|
| 5  | <u>Streptomyces</u>      |   |
|    | <u>caespitosus</u>       | mitomycins A, B, and C                              |
|    | <u>coelicolor</u>        | actinorhodin  |
| 10 | <u>coeruleorubidicus</u> | daunomycin  |
|    | <u>cyaneus</u>           | ditrisarubicin                                      |
|    | <u>flavogriseus</u>      | cyanocycline A                                      |
|    | <u>galilaeus</u>         | sclacinomycin A,<br>auramycins, and<br>sulfurmycins |
| 15 | <u>lusitanus</u>         | naphthyridinomycin                                  |
|    | <u>peuceticus</u>        | daunomycin and<br>adriamycin                        |
|    | <u>violochromogenes</u>  | arugomycin  |
|    | <u>E</u>                 |   |
| 20 | <u>cattleys</u>          | MM 4550, and MM 13902                               |
|    | <u>chartreusis</u>       | thienamycin<br>SF 1623 and<br>cephamycin A and B    |

X-6808A

-14-

TABLE IV $\beta$ -Lactam Antibiotic-Producing Organisms

|    | <u>Organism</u>  | <u>Antibiotic</u>   |
|----|--|---|
| 5  | <u>Agrobacterium</u>   | various $\beta$ -lactams  |
| 10 | <u>Cephalosporium</u><br><u>acremonium</u>                   | penicillins and<br>cephalosporins                                   |
|    | <u>Chromobacterium</u>                                       | various $\beta$ -lactams  |
| 15 | <u>Gluconobacter</u>   | various $\beta$ -lactams  |
|    | <u>Nocardia</u><br><u>lactamadurans</u><br><u>uniformis</u>  | cephamycin C<br>nocardicin  |
| 20 | <u>Penicillium</u><br><u>chrysogenum</u>                     | various penicillins and<br>other $\beta$ -lactams                   |
| 25 | <u>Serratia</u><br><u>antibioticus</u><br><u>argenteolus</u> | various $\beta$ -lactams<br><br>clavulanic acid<br>asparenomycin A, |

2233

87265660

X-6808A

-15-

TABLE IV (Continued)

|    | <u>Organism</u>          | <u>Antibiotic</u>   |
|----|--------------------------|---|
| 5  | <u>Streptomyces</u>      |   |
|    | <u>cinnamensis</u>       | cephamycin A and B  |
|    | <u>clavuligerus</u>      | PA-32413-I, cephamycin C,<br>A16886A, penicillins<br>cephalosporins,  |
| 10 | clavulanic               | acid, and other clavams   |
|    | <u>fimbriatus</u>        | cephamycin A and B  |
|    | <u>flavovirens</u>       | MM 4550 and MM 13902  |
|    | <u>flavus</u>            | MM 4550 and MM 13902  |
| 15 | <u>fulvoviridis</u>      | MM 4550 and MM 13902  |
|    | <u>griseus</u>           | cephamycin A and B<br>and carpetimycin A and B  |
|    | <u>halstedii</u>         | cephamycin A and B  |
|    | <u>heteromorphus</u>     | C2081X and  |
| 20 | <u>hygroscopicus</u>     | cephamycin A and B  |
|    | <u>lipmanii</u>          | deacetoxycephalosporin C<br>cephamycin, penicillin N,<br>7-methoxycephalosporin C,<br>A16884, MM4550, MM13902 |
| 25 | <u>olivaceus</u>         | epithienamycin F,<br>MM 4550, and MM 13902  |
|    | <u>panayensis</u>        | C2081X and<br>cephamycin A and B  |
|    | <u>rochei</u>            | cephamycin A and B  |
| 30 | <u>sioyaensis</u>        | MM 4550 and MM 13902  |
|    | sp. OA-6129              | OA-6129A  |
|    | sp. KC-6643              | carpetimycin A  |
|    | <u>virid chromogenes</u> | cephamycin A and B  |
|    | <u>wadayamensis</u>      | WS-3442-D   |

35

2234

87265660



X-6808A

-16-

TABLE V

Macrolide, Lincosamide, and Streptogramin  
Antibiotic-Producing Organisms

|    |                          |                                 |
|----|--------------------------|---------------------------------|
| 5  | <u>Organism</u>          | <u>Antibiotic</u>               |
|    | <u>Micromonospora</u>    |                                 |
|    | <u>rosaria</u>           | rosaramicin                     |
| 10 | <u>Streptomyces</u>      |                                 |
|    | <u>albireticuli</u>      | carbomycin                      |
|    | <u>albogriseolus</u>     | mikonomycin                     |
|    | <u>albus</u>             | albomycetin                     |
| 15 | <u>albus</u> var.        |                                 |
|    | <u>coilmyceticus</u>     | coleimycin                      |
|    | <u>ambofaciens</u>       | spiramycin and<br>foromacidin D |
|    | <u>antibioticus</u>      | oleandomycin                    |
| 20 | <u>avermittilis</u>      | avermectins                     |
|    | <u>bikiniensis</u>       | chalconomycin                   |
|    | <u>brunaeogriseus</u>    | albocycline                     |
|    | <u>caelestis</u>         | M188 and celesticetin           |
|    | <u>cinerochromogenes</u> | cineromycin B                   |
| 25 | <u>cirratus</u>          | cirramycin                      |
|    | <u>deltae</u>            | deltamycins                     |
|    | <u>djakartensis</u>      | niddamycin                      |
|    | <u>erythreus</u>         | erythromycins                   |
|    | <u>euroidicus</u>        | methymycin                      |
| 30 | <u>eurythermus</u>       | angolamycin                     |
|    | <u>fasciculus</u>        | amaromycin                      |

X-6808A

-17-

TABLE V (Continued)

|    | <u>Organism</u>                      | <u>Antibiotic</u>                          |
|----|--------------------------------------|--|
| 5  | <u>Streptomyces</u>                  |  |
|    | <u>felleus</u>                       | argomycin and<br>picromycin                |
|    | <u>fimbriatus</u>                    | amaromycin                                 |
|    | <u>flavochromogenes</u>              | amaromycin and<br>shincomycins             |
| 10 | <u>fradiae</u>                       | tylosin                                    |
|    | <u>fungicidicus</u>                  | NA-181                                     |
|    | <u>fungicidicus</u> var.             |  |
|    | <u>espinomyceticus</u>               | espinomycins                               |
| 15 | <u>furdicidicus</u>                  | mydecamycin                                |
|    | <u>goshikiensis</u>                  | bandamycin                                 |
|    | <u>griseofaciens</u>                 | PA133A and B                               |
|    | <u>griseoflavus</u>                  | acumycin                                   |
|    | <u>griseofuscus</u>                  | bundlin                                    |
| 20 | <u>griseolus</u>                     | griseomycin                                |
|    | <u>griseospiralis</u>                | relomycin                                  |
|    | <u>griseus</u>                       | borrelidin                                 |
|    | <u>griseus</u> ssp. <u>sulphurus</u> | bafilomycins                               |
|    | <u>halstedii</u>                     | carbomycin and leucanacidin                |
| 25 | <u>hygroscopicus</u>                 | tylosin                                    |
|    | <u>hygroscopicus</u> subsp.          |  |
|    | <u>aureolacrimosus</u>               | milbemycins                                |
|    | <u>kitastoensis</u>                  | leucomycin A <sub>3</sub> and<br>josamycin |
| 30 | <u>lavendulae</u>                    | aldgamycin                                 |
|    | <u>lincolnensis</u>                  | lincomycin                                 |

X-6808A

-18-

TABLE V (Continued)

|    | <u>Organism</u>  | <u>Antibiotic</u>  |
|----|--|--|
| 5  | <u>loidensis</u><br><u>macrosporeus</u><br><u>maizeus</u><br><u>mycarofaciens</u>                                  | vernAMYcin A and B<br>carbAMYcin<br>ingramycin<br>acetyl-leukomycin,<br>and espinomycin    |
| 10 | <u>narbonensis</u><br><br><u>narbonensis</u> var.<br><u>josamyceticus</u>  | josamycin and<br>narbomycin<br><br>leucomycin A <sub>3</sub><br>and josamycin              |
| 15 | <u>olivochromogenes</u><br><u>platensis</u><br><u>rimosus</u><br><br><u>rochei</u>                                 | oleandomycin<br>platenomycin<br>tylosin and<br>neutramycin<br>lankacidin and<br>borrelidin |
| 20 | <u>rochei</u> var.<br><u>volubilis</u><br><u>roseochromogenes</u><br><u>roseocitreus</u>                           | T2636<br>albocycline<br>albocycline  |
| 25 | <u>spinichromogenes</u> var.<br><u>suragaoensis</u><br><u>tendae</u><br><u>thermotolerans</u><br><u>venezuelae</u> | kujimycins<br>carbomycin<br>carbomycin<br>methymycins                                      |
| 30 | <u>violaceoniger</u>   | lankacidins and<br>lankamycin  |

X-6808A

-19-

TABLE VIMiscellaneous Antibiotic-Producing Streptomyces

| 5  | <u>Antibiotic Type</u>           | <u>Streptomyces Species</u>  | <u>Antibiotic</u>   |
|----|----------------------------------|--|---|
|    | amino acid<br>analogues          | sp.  | cycloserine   |
| 10 | cyclopentane ring-<br>containing | <u>coelicolor</u><br><u>erythrochromogenes</u><br><u>kasugaensis</u> | methylenomycin A<br>sarkomycin<br>aureothricin and<br>thiolutin   |
| 15 |                                  | <u>violaceoruber</u>   | methylenomycin A  |
|    | nitro-containing                 | <u>venezuelae</u>  | chloramphenicol   |
| 20 | polyenes                         | <u>griseus</u><br><u>nodosus</u><br><u>noursei</u>                   | candicidin<br>amphotericin B<br>nystatin  |
| 25 | tetracyclines                    | <u>aureofaciens</u><br><br><u>rimosus</u>                            | tetracycline, chlor<br>tetracycline,<br>demethyltetra<br>cycline, and<br>demethylchlortetra<br>cycline<br>oxytetracycline |

X-6808A

-20-

TABLE VII

## Nucleoside Antibiotic-Producing Organisms

|    | <u>Organism</u>  | <u>Antibiotic</u>  |
|----|--|--|
| 5  | <u>Corynebacterium</u><br><u>michiganese</u> pv. <u>rathayi</u>  | tunicamycin analogues  |
| 10 | <u>Nocardia</u><br><u>candidus</u>   | pyrazofurin  |
| 15 | <u>Streptomyces</u><br><u>antibioticus</u><br><u>chartreusis</u><br><u>griseoflavus</u> var.<br><u>thuringiensis</u><br><u>griseolus</u><br><u>lysosuperificus</u> | ara-A<br>tunicamycin<br><br>streptoviridans<br>sinefungin<br>tunicamycin |

X-6808A

-21-

TABLE VIII

## Peptide Antibiotic-Producing Organisms

| 5  | <u>Organism</u>         | <u>Antibiotic</u>                      |
|----|-------------------------|--|
|    | <u>Actinoplanes</u>     |  |
|    | <u>missouriensis</u>    | actaplanin                             |
|    | <u>teichomyceticus</u>  | teicoplanin                            |
| 10 | <u>Bacillus</u>         |  |
|    | various species         | bacitracin, polymixin,<br>and colistin |
| 15 | <u>Nocardia</u>         |  |
|    | <u>candidus</u>         | A-35512 and avoparcin                  |
|    | <u>lurida</u>           | ristocetin                             |
|    | <u>orientalis</u>       | vancomycin                             |
| 20 | <u>Streptomyces</u>     |  |
|    | <u>antibioticus</u>     | actinomycin                            |
|    | <u>aureus</u>           | thiostrepton                           |
|    | <u>canus</u>            | amphotycin                             |
|    | <u>eburosporeus</u>     | LL-AM374                               |
| 25 | <u>haranomachiensis</u> | vancomycin                             |
|    | <u>pristinaspinalis</u> | pristinamycin                          |
|    | <u>roseosporus</u>      | lipopeptides, such as<br>A21978C       |
|    | <u>toyocaensis</u>      | A47934                                 |
| 30 | <u>virginiae</u>        | A41030                                 |

X-6808A

-22-

TABLE IX

## Polyether Antibiotic-Producing Organism

|    | <u>Organism</u>          | <u>Antibiotic</u>                       |
|----|--------------------------|---|
| 5  | <u>Actinomadura</u>      |   |
|    | various species          | various polyethers                      |
|    | <u>oligosporus</u>       | A80190                                  |
| 10 | <u>Dactylosporangium</u> |   |
|    | various species          | various polyethers                      |
|    | <u>Nocardia</u>          |   |
| 15 | various species          | various polyethers                      |
|    | <u>Streptomyces</u>      |   |
|    | <u>albus</u>             | A204, A28695A and B,<br>and salinomycin |
| 20 | <u>aureofaciens</u>      | narasin                                 |
|    | <u>bobili</u>            | A80438                                  |
|    | <u>cacaoi</u> var.       |   |
|    | <u>ascoensis</u>         | lysocellin                              |
|    | <u>chartreusis</u>       | A23187                                  |
| 25 | <u>cinnamomensis</u>     | monensin                                |
|    | <u>conglobatus</u>       | ionomycin                               |
|    | <u>eurocidicus</u> var.  |   |
|    | <u>asterocidicus</u>     | laidlomycin                             |
|    | <u>flaveolus</u>         | CP38936                                 |
| 30 | <u>gallinarius</u>       | RP 30504                                |
|    | <u>griseus</u>           | grisorixin                              |

X-6808A

-23-

TABLE IX continued

|    | <u>Organism</u>   | <u>Antibiotic</u>  |
|----|---|--|
| 5  | <u>hygroscopicus</u>  | A218, emeracid, DE3936,<br>A120A, A28695A and B,<br>etheromycin, and<br>dianemycin |
| 10 | <u>lasaliensis</u><br><u>longwoodensis</u><br><u>mutabilis</u><br><u>pactum</u><br><u>ribosidificus</u><br><u>violaceoniger</u> | lasalocid<br>lysocellin<br>S-11743a<br>A80438<br>lonomycin<br>nigericin            |
| 15 | <u>Streptoverticillium</u><br>various species   | polyethers   |

20           The present invention is best exemplified by  
transforming antibiotic-producing microorganisms with  
genes that code for enzymes that catalyze chemical  
reactions governing the conversion of primary metabo-  
25   Q-methyltransferase, catalyzes the final step in the  
biosynthesis of tylosin. Transforming tylosin-producing  
microorganisms with a macrocin Q-methyltransferase-encoding



gene, designated as tylF, results in an improved tylosin biosynthetic pathway because observed are increased levels of the tylF gene product in the transformed cells.

5                   Accordingly, the present invention also provides a method for increasing the tylosin or tylosin precursor-producing ability of a tylosin-producing micro-organism, which comprises culturing a microorganism which produces tylosin or a tylosin precursor by a  
10   biosynthetic pathway, said microorganism being transformed with a DNA cloning vector or portion thereof which contains tylosin or tylosin-precursor biosynthetic gene coding for expression of a rate-limiting enzyme or  
15   gene product of the biosynthetic pathway, under conditions suitable for cell growth, expression of the tylosin or tylosin-precursor biosynthetic gene and production of the tylosin or tylosin precursor, provided that the culturing process provides an increase in the  
20   tylosin- or tylosin precursor-producing ability of the microorganism.

                  The present invention utilizes antibiotic biosynthetic genes to increase the antibiotic-producing ability of an organism. A small number of antibiotic biosynthetic genes have been cloned, characterized, and  
25   described in the relevant literature. Methods for isolating antibiotic biosynthetic genes have been developed, but one especially preferred method is described in Baltz et al., U.S. Patent Application serial number 742,349, filed June 7, 1985, (equivalent  
30   to European Pat. Appl. No. 86304239.6, Pub. No. 204,549) which is incorporated by reference. The present tylosin

antibiotic biosynthetic genes used in a specific exemplification of the present method initially were isolated from a  $\lambda$  library constructed in substantial accordance with the procedure described in Fishman et al., 1985, J. Bacteriology 161:199-206.

5 A schematic representation of the tylosin biosynthetic pathway is presented in Figure 1; each arrow in Figure 1 represents a step which is catalyzed by one or more tylosin biosynthetic gene products. The gene(s) responsible for each conversion is indicated above each arrow. Each genotypic designation may represent a class of genes that contribute to the same phenotype. A number of expression vectors are used to exemplify the present invention. These vectors comprise one or more tylosin biosynthetic genes and can be obtained from the Northern Regional Research Laboratories (NRRL), Peoria, Illinois 61604. Table X provides a brief description of each of the plasmids used to exemplify the method of the present invention.

20

Table X

## Plasmids Comprising Tylosin Biosynthetic Genes

| <u>H st/Designation</u>          | <u>Tylosin Gene(s)</u> | <u>NRRL</u> | <u>Date of Deposit</u> | <u>Map</u> |
|----------------------------------|------------------------|-------------|------------------------|------------|
| <u>E. coli</u> K12 HB101/pHJL280 | D, E, F, H, J          | B-18043     | February 18, 1986      | Fig. 2     |
| <u>E. coli</u> K12 HB101/pHJL284 | C, F, J                | B-18044     | February 18, 1986      | Fig. 3     |
| <u>E. coli</u> K12 HB101/pHJL309 | L, M                   | B-18045     | February 18, 1986      | Fig. 4     |
| <u>E. coli</u> K12 HB101/pHJL311 | C, F, J, K, H          | B-18046     | February 18, 1986      | Fig. 5     |
| <u>E. coli</u> K12 JM109/pHJL315 | D, E, F, H, J          | B-18047     | February 18, 1986      | Fig. 6     |

A number of Streptomyces fradiae strains are described which have mutant tylosin biosynthetic genes and, therefore, make much less tylosin than the strain from which they were derived. Table XI provides a brief  
5 description of these mutant strains.

Table XI

Streptomyces fradiae Mutants Defective in  
Tylosin Biosynthesis

| <u>Strain Designation</u> | <u>Mutant Gene</u> | <u>ATCC* or NRRL Accession No.</u>    |
|---------------------------|--------------------|---------------------------------------|
| GS15                      | <u>tylF</u>        | NRRL 18058 (deposited March 19, 1986) |
| GS16                      | <u>tylE</u>        | ATCC 31664 (publicly available)       |
| GS28                      | <u>tylF</u>        | NRRL 18059 (deposited March 19, 1986) |
| GS33                      | <u>tylL</u>        |                                       |
| GS48                      | <u>tylD</u>        | NRRL 12170 (publicly available)       |
| GS52                      | <u>tylC</u>        | NRRL 18060 (deposited March 19, 1986) |
| GS62                      | <u>tylM</u>        |                                       |
| GS76                      | <u>tylD</u>        |                                       |
|                           | <u>tylH</u>        | NRRL 12171 (publicly available)       |
| GS85                      | <u>tylK</u>        |                                       |
| GS88                      | <u>tylJ</u>        |                                       |

\*ATCC is the American Type Culture Collection, Rockville, MD 20852,  
and NRRL is the Northern Regional Research Laboratory, Peoria, IL 61604.

Plasmids pHJL280, pHJL284, and pHJL315 were used to transform Streptomyces fradiae GS15 and Streptomyces fradiae GS28. The GS15 and GS28 strains were prepared from S. fradiae C4 by nitrosoguanidine mutagenesis. S. fradiae C4 was derived from S. fradiae T59235 (ATCC 19609) by mutagenesis. The GS15 strain makes almost no tylosin, and the GS28 strain makes low levels of tylosin, as compared with the C4 strain. The decreased or nonexistent tylosin-producing ability of the GS15 and GS28 strains is believed to result from mutations affecting the tylF gene, which encodes macrocin O-methyltransferase (MOMT). The MOMT enzyme, which is required for the conversion of macrocin to tylosin in the tylosin biosynthetic pathway, is frequently present in reaction rate-limiting amounts in tylosin-producing strains. Plasmids pHJL280, pHJL284, and pHJL315 remove this reaction limitation by providing a means for increasing both the copy number of the tylF biosynthetic gene and also the concentration of macrocin O-methyltransferase available for tylosin biosynthesis. Accordingly, fermentation of S. fradiae GS15/pHJL280, S. fradiae GS15/pHJL284, S. fradiae GS15/pHJL315, S. fradiae GS28/pHJL284, S. fradiae GS28/pHJL280, and S. fradiae GS28/pHJL315 for 72 hours results in about a 2-fold to a 6-fold increase in the production of macrocin O-methyltransferase over that produced in the C4 strain and a 120-fold increase over that produced in the GS28 strain.

Plasmid pHJL280 was also used to transform:  
(1) Streptomyces fradiae GS16; (2) S. fradiae GS48; (3)

S. fradiae GS76; and (4) S. fradiae GS88 which produce tylosin below detection limits and were derived by mutagenesis of the C4 strain. Untransformed strains GS16, GS48, GS76, and GS88 respectively produce a

5 defective enzyme or a rate-limiting amount of (1) the tylE, demethylmacrocin O-methyltransferase, enzyme; (2) the tylD enzyme, which is required for addition or biosynthesis of 6-deoxy-D-allose; (3) the tylH enzyme,

10 which is required for oxidation of the C-23 methyl position of tylactone; and (4) the tylJ enzyme. Untransformed strains GS16, GS48, GS76, and GS88, respectively, tend to accumulate demethylmacrocin, demycinosyl tylosin, 23-deoxydemycinosyl tylosin, and demycinosyl tylosin rather than the desired tylosin

15 antibiotic compound.

Plasmid pHJL280 provides a means for increasing the efficiency of the tylosin biosynthetic pathway by not only providing a non-defective gene but also by increasing the copy number of the tylD, tylE, tylH, and

20 tylJ biosynthetic genes and by increasing the intracellular amount of the products specified by these genes. The concentration of available tylE gene product, therefore, is increased, resulting in an elevated amount of enzyme capable of driving the conversion of

25 demethylmacrocin to macrocin to tylosin in the tylosin biosynthetic pathway. Similarly, the concentration of available tylD, tylH, and tylJ gene products is also increased, resulting in the production of elevated amounts of the enzymes capable of driving the 6-deoxy-

30 D-allose addition and C-23 oxidation of tylosin

precursors. Fermentation of Streptomyces fradiae GS16/pHJL280, S. fradiae GS48/pHJL280, S. fradiae GS76/pHJL280, and S. fradiae GS88/pHJL280 for 144-168 hours results in yields of tylosin that are significantly increased over that of the untransformed, low-tylosin-producing, mutant strains. Such transformed strains have higher enzyme levels of the particular enzymes encoded on plasmid pHJL280 than the parent C4 strain and thus further exemplify the present invention. Plasmid pHJL280 can be used to improve the tylosin-producing ability of any organism in which the tylD, tylE, tylF, tylH, or tylJ gene products (or any combination thereof) are present in rate-limiting amounts for tylosin biosynthesis.

Plasmid pHJL284 was also used to transform Streptomyces fradiae GS52, a low tylosin-producing, mutant strain derived from the C4 strain that produces reaction-limiting amounts of an enzyme required for the biosynthesis or addition of mycarose to de-O-methyl-lactenocin. Thus, the tylosin biosynthetic pathway of Streptomyces fradiae GS52 tends to produce desmycosin rather than the desired tylosin antibiotic compound. Plasmid pHJL284 provides a means for improving the synthetic efficiency of this pathway by providing a non-defective biosynthetic gene and by increasing the copy number of the tylC biosynthetic gene. The concentration of available tylC gene product in the transformed strain, therefore, is increased, resulting in the elevated production of enzyme capable of driving the desired addition reaction. Accordingly, fermentation of



Streptomyces fradiae GS52/pHJL284 for 144-168 hours results in a level of tylosin production that is significantly increased over that of the untransformed mutant strain and results in higher tylC enzyme levels than those in the parent C4 strain. Plasmid pHJL284 was also used in the present method to improve the tylosin-producing ability of Streptomyces fradiae GS88, a tylJ mutant, and thus can also be used in the present method to improve the tylosin-producing ability of any organism in which the tylC, tylF, or tylJ gene products (or any combination thereof) are present in rate-limiting amounts for tylosin biosynthesis.

Plasmid pHJL309 contains the tylL and tylM biosynthetic genes and was used in the present method to improve the tylosin-producing ability of Streptomyces fradiae GS33, a tylL mutant, and GS62, a tylM mutant. Plasmid pHJL309 can also be used in the present method to improve the tylosin-producing ability of any organism in which the tylL or tylM gene products (or both) are present in rate-limiting amounts for tylosin biosynthesis.

Plasmid pHJL311 contains the tylC, tylF, tylH, tylJ, and tylK biosynthetic genes and so was used in the present method to improve the tylosin-producing ability of Streptomyces fradiae GS52, a tylC mutant; GS88, a tylJ mutant; GS15 and GS28, both of which are tylF mutants; and GS85, a tylK mutant. Plasmid pHJL311 can also be used in the present method to improve the tylosin-producing ability of any organism in which the tylC, tylF, tylH, tylJ, or tylK gene products (or any

combination thereof) are present in rate-limiting amounts for tylosin biosynthesis.

Plasmid pHJL315 contains the tylD, tylE, tylF, tylH, and tylJ biosynthetic genes and so was used in the present method to improve the tylosin-producing ability of Streptomyces fradiae GS48, a tylD mutant; GS88, a tylJ mutant; GS16, a tylE mutant; GS76, a tylD, tylH double mutant; and GS15 and GS28, both of which are tylF mutants. Plasmid pHJL315 can also be used in the present method to improve the tylosin-producing ability of any organism in which the tylD, tylE, tylF, tylH, or tylJ gene products (or any combination thereof) are present in rate-limiting amounts for tylosin biosynthesis.

These results demonstrate that the vectors of the present invention can increase the antibiotic-producing ability of an antibiotic-producing organism by providing higher enzyme or other gene product levels, as compared to an untransformed organism, of an enzyme or other gene product that is rate-limiting in an antibiotic biosynthetic pathway. However, plasmid maintenance in an antibiotic-producing host cell sometimes requires significant expenditures of the cell's energy, energy that might otherwise be used to produce antibiotic. Thus, certain microorganisms transformed with autonomously replicating vectors actually show a decrease in antibiotic-producing ability, even though the same vectors can increase the antibiotic-producing ability of other organisms. Not wishing the present invention to

X-6808A

-34-

be bound or limited in any way by theory, this apparent anomaly can be explained by the fact that antibiotics are produced from primary metabolites, such as acetate, propionate, malonyl-CoA, methylmalonyl-CoA, and glucose, by the action of specific enzymes. These enzymes are usually not present during the rapid growth phase of an organism and so do not rob the growing cell of needed compounds. As growth becomes limited by nutritional conditions, antibiotic biosynthetic genes are believed to be activated, causing the synthesis of enzymes that redirect the flow of certain primary metabolites into antibiotic products.

The synthesis of antibiotics is also believed to be a dispensable function in antibiotic-producing organisms because mutants blocked in the biosynthesis of antibiotics are viable and grow as well as the antibiotic-producing parent. Wild-type strains produce a relatively small amount of antibiotic, which is apparently adequate to provide the organism with a selective advantage.

The development of industrial antibiotic producing strains from natural isolates involves many cycles of mutation and selection for higher antibiotic production. Because the synthesis of antibiotics drains primary metabolites and cellular energy away from growth and maintenance functions, it is believed that selection for higher antibiotic production frequently occurs at the expense of the vitality of the organism. Thus, the generation of high antibiotic-producing strains involves finely balancing the cells nutritional and energy

resources between growth-maintenance functions and antibiotic production. As a consequence of this fine-tuning, high-yielding production strains tend to be extremely sensitive to factors that affect cellular physiology. For example, introduction of autonomously-replicating vectors, notably multicopy plasmids, sometimes tends to decrease the antibiotic-producing ability of an organism that normally produces antibiotics at high levels. The mechanism of this inhibition is not clear, but may occur at an early step in the biosynthesis of the antibiotic because measurable levels of antibiotic precursors do not accumulate under these conditions. In addition, autonomously replicating vectors may drain pools of precursors for DNA or RNA synthesis or, in high copy number, may titrate DNA binding proteins, such as RNA polymerase, DNA polymerase, polymerase activators, or repressors of gene expression. Another frequent limitation of autonomously replicating vectors is spontaneous loss. Spontaneous loss is especially problematical when the vector reduces growth rate as frequently occurs. Selection for a resistance marker on the plasmid can ensure the growth of homogeneous, plasmid-containing populations but can also disrupt the fine physiological balance (already mentioned) of an antibiotic fermentation. Selection for unstable plasmids operates by killing or inhibiting the bacteria that lose the plasmid and can result in a reduced growth rate.

The negative effect, sometimes observed, of autonomously replicating vectors on the

antibiotic-producing ability of a microorganism is  
greatest in high-producing strains that are delicately  
balanced with respect to growth-maintenance functions  
and antibiotic production. The present invention  
5 overcomes this previously unrecognized problem of the  
negative effect of autonomous plasmid replication on  
high-producing strains by providing methods of culturing  
the transformed host cell to facilitate identification  
of transformed cells containing integrated plasmid and,  
10 in addition, by providing vectors with features that  
also facilitate detection of integration. Selecting a  
culturing procedure that results in integration is  
important in improving the antibiotic-producing ability  
of highly selected and conventionally improved  
15 antibiotic-producing organisms. Organisms or strains  
that have a low antibiotic-producing ability can be  
improved by transformation via either integration or  
autonomous vector replication. As those skilled in the  
art of fermentation technology will appreciate, the  
20 greatest improvement in antibiotic-producing ability is  
shown when the present invention is applied to low  
antibiotic-producing strains.

(X) Integration of plasmid DNA is readily accom-  
plished by transforming a given antibiotic-producing  
25 strain or mutant thereof according to standard transfor-  
mation procedures, selecting or otherwise identifying  
the transformants, and then culturing the cells under  
conditions that do not require the presence of plasmid  
DNA sequences for the host cell to grow and replicate.  
30 After several generations under non-selective

conditions, certain cells will no longer contain free plasmid DNA, so by selecting for or otherwise identifying plasmid DNA sequences present in the host cell, one can identify host cells in which the plasmid DNA has integrated into the chromosomal (genomic) DNA of the cell. This culturing technique to obtain integration of vector DNA is especially useful when used in conjunction with a vector that is inherently unstable in the transformed host cell, so that culturing without selective pressure to maintain the vector generates segregants that are free of the plasmid. Bibb *et al.*, 1980, Nature 384:526-531, describe a DNA sequence needed for stable inheritance of a vector, and a variety of vectors have been constructed that lack this stability sequence.

For instance, cloning vectors pHJL210 and pHJL401, which were used to construct the plasmids of the present invention, lack this stability sequence. Plasmid pHJL210 is disclosed in U.S. Patent Application Serial No. 639,566, filed August 10, 1984, (equivalent to European Publication No. 176199). Plasmid pHJL401 is disclosed in U.S. Patent Application Serial No. 841,920, filed March 20, 1986, which is a continuation-in-part of Serial No. 763,172, filed August 7, 1985, (equivalent to European Application No. 86306011.7, filed August 5, 1986, Pub. 213,779). As used, "unstable" refers to plasmids that are lost at high frequency by transformed cells only when those cells are cultured in the absence of selective pressure for plasmid maintenance because, for example, plasmids such as pHJL210 and pHJL401 are quite stable

when selective pressure is applied to the transformed host cell. When host cells transformed with stable vectors are cultured in the absence of selective pressure, the vector is not lost with the high frequency observed with unstable vectors, and identification of integrants is made difficult by the great number of cells that still contain autonomously replicating plasmid even after growth under nonselective conditions. Selection for integrants is more fully described below. Once the vector DNA has integrated into the chromosomal DNA of the host cell, one observes the maximum increase in antibiotic-producing ability for that host cell, because inhibition by autonomously replicating plasmid no longer occurs.

Integration of vectors containing cloned genes into the genome of the producing organism can be achieved in a number of ways. One way is to use a lysogenic bacteriophage or other phage vector that can integrate into the genome of the host strain. Another approach is to use a plasmid vector carrying the cloned genes and to screen for integration of the recombinant plasmid into the host genome by a single recombination event between the cloned sequence and the homologous chromosomal sequence. Integration frequency of a vector can be dramatically increased by adding DNA homologous to the genomic DNA of the host cell to the vector. As used "integration" refers both to a single recombination event, known as Campbell-type recombination, and also to a double-crossover event, which results in exchange of genetic information between the vector and the

chromosome. With double-crossover recombination, only a portion of the vector integrates into the chromosomal DNA.

For example, a plasmid carrying cloned tylosin biosynthetic genes (tyl) could integrate into the Streptomyces fradiae genome by a single crossover between the tyl genes on the plasmid and the homologous tyl genes in the genome. Another option would be to put a non-tyl S. fradiae DNA sequence on the plasmid in addition to the cloned tyl genes and to screen for integration at the locus corresponding to the non-tyl sequence. The latter approach avoids the possible mutagenic effects of integration into the tyl sequences, but if double-crossover recombination is desired, the vector should comprise the antibiotic biosynthetic genes flanked by separate sequences of homologous DNA.

To avoid the potentially adverse effects, however remote, of a recombinant plasmid (either autonomously replicating or integrated) on tylosin production, one can make use of the ability of Streptomyces fradiae to take up tylosin precursors from the culture medium and convert them to tylosin. In one fermentation of a tylosin-producing strain that had been transformed with plasmid pHJL280 and cultured to obtain integrants, only a subpopulation (~18%) of the cells were thiostrepton resistant, which indicates the presence of plasmid pHJL280 sequences. However, this subpopulation contained multiple copies of the genes for two rate-limiting enzymes, demethylmacrocin-O-methyltransferase (DMOMT) and macrocin-O-methyltransferase (MOMT), and



consequently elevated (about 9 fold) levels of the two enzymes, and was able to convert all of the normally accumulated demethylmacrocin and macrocin to tylosin (see Table XIV).

5           Thus, one can develop specific strains of S. fradiae containing multiple copies of rate-limiting genes and high enzyme levels to act as converters of accumulated precursors to tylosin. These converter strains can be used in several different ways: (1) the  
10 converter strain can be co-inoculated into the fermentor with the normal production strain at a low ratio of converter:producer; (2) the converter strain can be introduced into a production fermentation culture late in the cycle to convert intermediates; (3) the converter  
15 strain can be kept in a separate "reactor", to which the fermentation production broth from the producer strain would be added; or (4) the converter strain can be immobilized on a column, and fermentation broth from the  
20 producer strain passed through. Those skilled in the art will recognize that having separate production and converting populations eliminates the adverse effects that recombinant plasmids sometimes have on antibiotic production in high antibiotic-producing strains.

25           Separate populations also eliminate vector stability problems, because the converting strains can be grown in small vessels in which antibiotic selection or some other selection means for maintenance of the plasmid can be carefully regulated and controlled. In  
30 essence, the converting strain is a source of enzymes, and the production of these enzymes at high level can be

X-6808A

-41-

approached in much the same way as production of proteins from recombinant plasmids in E. coli.

Of course, antibiotic production is only increased by the method of the present invention when the transforming DNA comprises a gene that encodes the rate-limiting enzyme of the untransformed strain. Various methods for determining the rate-limiting step in the biosynthesis of an antibiotic are known in the art (Seno and Baltz, 1982, Antimicrobial Agents and Chemotherapy 21:758-763), but there is no need to identify the rate-limiting step when the entire set of antibiotic biosynthetic genes are available for introduction into the antibiotic-producing strain. If a rate-limiting enzyme is not known, the antibiotic-producing strain is transformed with the entire set of antibiotic biosynthetic genes, thus ensuring that, no matter what enzyme is rate-limiting, the transformed host cell will have higher levels of the rate-limiting enzyme than the untransformed host cell. Often, however, the rate-limiting enzyme of an antibiotic biosynthesis pathway will be known, and the method of the present invention can be used to increase the antibiotic-producing ability of the organism by transforming the organism with a vector that encodes the rate-limiting antibiotic biosynthetic enzyme.

For instance, the GS15 strain, which produces no readily detectable tylosin (the level of tylosin produced by these cells is below the detection limits for the assay used to determine tylosin levels) and the GS28 strain, which produces very low levels of tylosin,

contain tylF mutations, so that it is a relatively simple matter to identify the rate-limiting step in tylosin biosynthesis in these mutant strains. The strain from which the GS15 and GS28 strains were derived, designated Streptomyces fradiae C4, produces high levels of tylosin and accumulates relatively large amounts of macrocin, the immediate precursor of tylosin on which the tylF gene product acts to form tylosin. Other S. fradiae strains that produce even more tylosin than the C4 strain accumulate even more macrocin than the C4 strain. These observations indicate that the tylF gene product is present in rate-limiting amounts for the biosynthesis of tylosin in high tylosin-producing strains. Transformation of these macrocin-accumulating strains with a vector comprising the tylF gene followed by isolation of those transformants that only contain integrated copies of the vector yields transformants that produce more tylosin than the untransformed cells. The increase in tylosin production observed in these transformants is related to the amount of macrocin that accumulates in the untransformed cells. It will be apparent to those skilled in the art that the transformants produced by the foregoing procedure might still contain rate-limiting amounts of the tylF gene product, in which case a further increase of the tylF copy number would further increase tylosin yield, or the transformed strains might now contain rate-limiting amounts of yet another antibiotic biosynthetic enzyme, the level of

which could be made non-rate-limiting by the method of the present invention.

The present invention provides both a method and recombinant DNA cloning vectors for increasing the production of an antibiotic by manipulation of antibiotic biosynthetic pathways. An illustrative antibiotic biosynthetic pathway involves the biosynthesis of tylosin, a complex macrolide produced by strains of Streptomyces fradiae, Streptomyces rimosus, and Streptomyces hygroscopicus. Tylosin is composed of a 16-member branched lactone (tylonolide) to which three sugars (mycarose, mycaminose, and mycinose) are attached. The lactone is derived from two acetates, five propionates, and a butyrate by condensation of a propionyl-S-coenzyme A molecule with two malonyl-S-coenzyme A molecules, four methylmalonyl-S-coenzyme A molecules, and an ethylmalonyl-S-coenzyme A molecule by a scheme believed analogous to that involved in fatty acid biosynthesis. Lactone formation, sugar biosynthesis/attachment, and the conversion of resultant intermediate compounds to tylosin are catalyzed by a series of gene-encoded enzymes. Cloning genes that code for such enzymes allows modification and improvement in the operational efficiency of the tylosin biosynthetic pathway and is illustrative of the present invention.

Illustrative tylosin biosynthetic genes that can be used for purposes of the present invention include, for example, the tylC, tylD, tylE, tylF, tylH, tylJ, tylK, tylL, and tylM, genes. Of this group, the tylF gene is preferred because the macrocin

O-methyltransferase enzyme encoded thereby appears to be rate-limiting in the tylosin biosynthetic pathway of most tylosin-producing strains. Macrocin accumulates to unacceptable levels under conditions of optimum fermentation of Streptomyces fradiae because of the rate-limiting steps catalyzed by the tylF gene product. The tylF enzyme catalyzes the conversion of macrocin to tylosin, as depicted in Figure 1 of the accompanying drawings. Over-production of the tylF gene product, macrocin O-methyltransferase, results in the more efficient operation of the tylosin biosynthetic pathway as indicated by increased antibiotic yield and lower cost of fermentation.

Those skilled in the art will recognize that the present invention is not limited to the use of plasmids pHJL280, pHJL284, pHJL309, pHJL311, or pHJL315. The antibiotic biosynthetic genes contained in these vectors can be excised in whole or in part and ligated into any number of different recombinant DNA cloning vectors. For instance, digestion of plasmid pHJL280 with restriction enzymes BamHI and BglII yields five BamHI-BamHI fragments with sizes of ~10.3 kb, ~6.54 kb, ~2.3 kb, ~1.7 kb, and ~1.0 kb; two BamHI-BglII fragments with sizes of ~2.9 kb and 2.0 kb; and one BglII-BglII fragment ~0.2 kb in size. The ~2.9 kb BamHI-BglII fragment of plasmid pHJL280 contains the tylF gene. Digestion of plasmid pHJL280 with restriction enzymes BglII and EcoRI generates four fragments: an ~11.24 kb EcoRI-EcoRI fragment; an ~11.5 kb BglII-EcoRI fragment; an ~4.0 kb EcoRI-BglII fragment, and an ~0.2 kb

BglII-BglII fragment. The ~4.0 kb EcoRI-BglII fragment of plasmid pHJL280 contains the tylE gene.

Digestion of plasmid pHJL284 with restriction enzymes BamHI and EcoRI generates three BamHI-BamHI fragments with sizes of ~9.7 kb, ~2.3 kb, and ~1.0 kb; and four EcoRI-BamHI fragments with sizes of ~6.24 kb, ~4.3 kb, ~2.3 kb, and ~1.1 kb. The ~2.3 kb BamHI-EcoRI fragment of plasmid pHJL284 contains the tylF gene. Digestion of plasmid pHJL284 with restriction enzyme EcoRI generates two fragments with sizes of ~1.4 kb and ~10.54 kb; the ~16.4 kb fragment contains the tylF, tylC, and tylJ genes. The ~1.7 kb EcoRI-BamHI restriction fragment of plasmid pHJL311 comprises the tylK gene. The ~18.5 kb EcoRI restriction fragment, as well as the ~8.3 kb BamHI-KpnI restriction fragment, of plasmid pHJL309 contains the tylL and tylM genes.

Any of the aforementioned tyl gene-containing fragments can be ligated into other vectors to make vectors useful in the present method. Such other vectors include, for example, those vectors disclosed in U.S. Patent Nos. 4,468,462; 4,513,086; 4,416,994; 4,503,155; and 4,513,185; and also plasmids pIJ101, pIJ350, pIJ702 (ATCC 39155), SCP2\* (NRRL 15041), pHJL192, pHJL197, pHJL198, pHJL210, pHJL211, pHJL400, pHJL401, pHJL302, pIJ922, pIJ903, pIJ941, pIJ940, and pIJ916. These vectors replicate in Streptomyces fradiae and other tylosin-producing strains and are thus useful for cloning the present antibiotic biosynthetic genes. The "unstable" vectors described above are preferred when integration of the vector is desired.

Illustrative Streptomyces strains that can be used for purposes of the present invention include, for example, S. fradiae, S. fradiae GS52, S. fradiae GS48, S. fradiae GS16, S. fradiae GS28, S. fradiae GS15, S. fradiae GS76, S. rimosus, and S. hygrosopicus. Streptomyces hygrosopicus and S. rimosus are well known, having been deposited at the American Type Culture Collection (ATCC), Rockville, Maryland 20852. A number of strains of S. hygrosopicus can be obtained under the accession numbers ATCC 27438, ATCC 21449, ATCC 15484, ATCC 19040, and ATCC 15420, and S. rimosus can be obtained under the accession number ATCC 10970. Of the Streptomyces taxa, S. fradiae GS16, S. fradiae GS15, and S. fradiae GS28 are preferred, especially for transformation with plasmid pHJL280. Streptomyces fradiae is also an especially well known microorganism and several strains are available, on an unrestricted basis, from the Northern Regional Research Laboratory (NRRL), Peoria, Illinois 61604 and the ATCC under the respective accession numbers NRRL 2702, NRRL 2703, and ATCC 19609.

The recombinant plasmids described in the present invention each comprise one or more antibiotic biosynthetic genes. Unless part of a polycistron, each antibiotic biosynthetic gene normally comprises: (1) a promoter that directs transcription of the gene; (2) a sequence that, when transcribed into mRNA, directs translation of the transcript; (3) a protein-coding sequence; and (4) a transcription terminator. Each of these elements is independently useful and can, through the techniques of recombinant DNA technology, be used to

form recombinant genes of great variety. As one example, the protein-coding sequence for the tylF gene can be linked to the promoter, translation-activating sequence, and transcription-terminating sequence from a non-Streptomyces fradiae gene to form a recombinant gene that functions in the host from which the non-S. fradiae sequences were isolated. Such a novel gene could be used to produce a hybrid antibiotic if introduced into an organism that produced an antibiotic or antibiotic intermediate that is not found in the tylosin pathway but that would serve as a substrate for the novel gene product. Similarly, the promoter and other regulatory elements of the tylF gene could be linked to the coding sequence of a non-tylosin antibiotic biosynthetic gene to prepare a hybrid gene that would function in S. fradiae. Thus, the individual elements of each of the antibiotic biosynthetic genes on each of the plasmids described herein comprise an important component of the present invention.

For example, sequence data on the tylF nucleotide sequence has identified the tylF promoter, which comprises an important aspect of the present invention. The sequence (only one strand of which is depicted for convenience) is shown below; the promoter and translation-activating sequence of the tylF gene are believed to reside in the sequence between residues 1



0238323

and 207. The sequence nds with the beginning of the coding region of the tylF gene.

As an additional embodiment, the gene sequence for the tylF gene is provided. In particular, the entire tylF gene, including the promoter and translation-activating sequence noted above, has been found to have the following sequence:

5

10 20 30 40

5'-TTC GCG GGA TGG ATG CTG ACC CGG GGG TCG GCC AGC AGC GCC CGG ACG

10 50 60 70 80 90

TGA TCT GGC GGG AGA TCA GCC AGA CCG GCG CCC CGT CCC ACA GCT CGG

100 110 120 130 140

CCC GGG CGA TCG GCT CCT CCG CCC GGA GGG CGG CGT ACT GCT CGG GAG

15 150 160 170 180 190

GGC TGA AGG GAC AGG TGC GGG CGA CCG GCC AGG CGA TGC TGC GCC GGC

200 210 220 230 240

20 CTG CGG CCC CGT CGG TGT CGT TGG CGC GTG CTG CGG GCA ACA GAA TCC

250 260 270 280

CCT TTT GTG ACG GGC GGG CGT CCC CGG ACG AGG ACA CGA CTC GCT GCG

25 290 300 310 320 330

GCC TCA ACG AAA ACA CCG TGT CCG GTG CCC AGG CCA CGA ACG GTG ACC

340 350 360 370 380

GGT CTG TGT CAG GTC GCC CGT GGT GAC GGG CTC CGG GGC TGC GGC GCG

TTCAG

TAAAT

2267

87265660

390 400 410 420 430  
 GGC GGC CGA CCT TGA CAT ACC CGC GGC CGG GCT CCT CGT TCC GGC GCG  
 5 440 450 460 470 480  
 GCC CGC GCC GAT AGC GTC CGT CCT CAC CGG CTC CGG CGT CCG CGT CCC  
 490 500 510 520  
 CGC CGG GAC GTG CCA CCT CTC CCG ACC CCG CGA GCC GAT CGA CCC GCT  
 10 530 540 550 560 570  
 ACT GGA GGA CCC GTG GCA CCT TCC CCG GAC CAC GCC CGC GAT CTC TAC  
 VAL ALA PRO SER PRO ASP HIS ALA ARG ASP LEU TYR  
 5 10  
 15 580 590 600 610 620  
 ATC GAG CTG CTG AAG AAG GTC GTC TCG AAC GTC ATC TAC GAG GAC CCC  
 ILE GLU LEU LEU LYS LYS VAL VAL SER ASN VAL ILE TYR GLU ASP PRO  
 15 20 25  
 20 630 640 650 660 670  
 ACC CAT GTG GCG GGG ATG ATC ACC GAC GCG TCG TTC GAC CGG ACG TCC  
 THR HIS VAL ALA GLY MET ILE THR ASP ALA SER PHE ASP ARG THR SER  
 30 35 40  
 25 680 690 700 710 720  
 CGT GAG AGC GGC GAG GAC TAC CCC ACG GTC GCC CAC ACG ATG ATC GGC  
 ARG GLU SER GLY GLU ASP TYR PRO THR VAL ALA HIS THR MET ILE GLY  
 45 50 55 60

0238323

-50-

5                   730                   740                   750                   760  
 CTC AAG CGT CTG GAC AAT CTC CAC CGG TGC CTC GCG GAC GTC GTG GAG  
 LEU LYS ARG LEU ASP ASN LEU HIS ARG CYS LEU ALA ASP VAL VAL GLU  
                   65                   70                   75

10                   770                   780                   790                   800                   810  
 GAC GGC GTC CCC GGT GAC TTC ATC GAG ACC GGG GTG TGC CGC GCG CCG  
 ASP GLY VAL PRO GLY ASP PHE ILE GLU THR GLY VAL CYS ARG ALA PRO  
                   80                   85                   90

15                   820                   830                   840                   850                   860  
 TGC ATC TTC GCC CGC GGA CTG CTG AAC GCG TAC GGC CAG GCC GAC CGC  
 CYS ILE PHE ALA ARG GLY LEU LEU ASN ALA TYR GLY GLN ALA ASP ARG  
                   95                   100                   105

20                   870                   880                   890                   900                   910  
 ACC GTC TGG GTC GCC GAC TCC TTC CAG GGC TTT CCC GAG CTG ACC GGG  
 THR VAL TRP VAL ALA ASP SER PHE GLN GLY PHE PRO GLU LEU THR GLY  
                   110                   115                   120

25                   920                   930                   940                   950                   960  
 TCC GAC CAC CCG CTG GAC GTC GAG ATC GAC CTC CAC CAG TAC AAC GAG  
 SER ASP HIS PRO LEU ASP VAL GLU ILE ASP LEU HIS GLN TYR ASN GLU  
                   125                   130                   135                   140

30                   970                   980                   990                   1000  
 GCC GTG GAC CTG CCC ACC AGC GAG GAG ACC GTG CGG GAG AAC TTC GCC  
 ALA VAL ASP LEU PRO THR SER GLU GLU THR VAL ARG GLU ASN PHE ALA  
                   145                   150                   155

0238323

-51-

|    |   |      |      |      |      |
|----|---|------|------|------|------|
|    | 1010  | 1020 | 1030 | 1040 | 1050 |
|    | CGG TAC GGG CTG CTC GAC GAC AAC GTC CGT TTC CTG GCG GGG TGG TTC |      |      |      |      |
|    | ARG TYR GLY LEU LEU ASP ASP ASN VAL ARG PHE LEU ALA GLY TRP PHE |      |      |      |      |
| 5  |   | 160  |      | 165  | 170  |
|    | 1060  | 1070 | 1080 | 1090 | 1100 |
|    | AAG GAC ACC ATG CCG GCT GCG CCC GTG AAG CAG CTC GCG GTG ATG CCG |      |      |      |      |
|    | LYS ASP THR MET PRO ALA ALA PRO VAL LYS GLN LEU ALA VAL MET ARG |      |      |      |      |
| 10 |   | 175  |      | 180  |      |
|    | 1110  | 1120 | 1130 | 1140 | 1150 |
|    | CTG GAC GGC GAC TCC TAC GGC GCC ACC ATG GAT GTG CTC GAC AGC CTG |      |      |      |      |
|    | LEU ASP GLY ASP SER TYR GLY ALA THR MET ASP VAL LEU ASP SER LEU |      |      |      |      |
| 15 |   | 190  |      | 195  | 200  |
|    | 1160  | 1170 | 1180 | 1190 | 1200 |
|    | TAC GAG CCG CTG TCG CCG GGC GGT TAC GTC ATC GTC GAC GAC TAC TGC |      |      |      |      |
|    | TYR GLU ARG LEU SER PRO GLY GLY TYR VAL ILE VAL ASP ASP TYR CYS |      |      |      |      |
| 20 |   | 205  |      | 210  | 215  |
|    | 1210  | 1220 | 1230 | 1240 |      |
|    | ATC CCG GCC TGC CGC GAG CCG TGC ACG ACT TCC GCG ACC GGC TCG GCA |      |      |      |      |
|    | ILE PRO ALA CYS ARG GLU ARG CYS THR THR SER ALA THR GLY SER ALA |      |      |      |      |
| 25 |   | 225  |      | 230  | 235  |
|    | 1250  | 1260 | 1270 | 1280 | 1290 |
|    | TCC GCG ACA CGA TCC ACC GGA TCG ACC GCC AGG GCG CTA TTG GCG GCA |      |      |      |      |
|    | SER ALA THR ARG SER THR GLY SER THR ALA ARG ALA LEU LEU ALA ALA |      |      |      |      |
| 30 |   | 240  |      | 245  | 250  |

2270

87265660

1300 1310 1320 1330 1340  
 CAG CGG CTG AGT CGT TCC GCC CGA GAG CCC GAC GAG AGC AGG AGA TAT  
 GLN ARG LEU SER ARG SER ALA ARG GLU PRO ASP GLU SER ARG ARG TYR  
 5 255 260 265  
  
 1350 1360 1370 1380 1390  
 GCG AGA CAC GAC GCG CCC GCT CGG CAT TGA GGG AGC GTG GGT GAT CCA  
 ALA ARG HIS ASP ALA PRO ALA ARG HIS  
 10 270 275  
  
 1400 1410 1420 1430 1440  
 GCC GGA GAT CCA TCC GGA CCG GCG CGG CGA GTT CCA CGC GTG GTT CCA  
 15  
 1450 1460 1470 1480  
 GAG CCA GCC GAG TTC CCG CCG CTG ACC GGT CAC TCC TTC TCC GTG CCG  
  
 1490 1500 1510 1520 1530  
 CAG GTC GTC AAT ATC GCG TGT CCC GGA AAG GCG CCG CTG CCG CAT CCA  
 20  
 1540 1550 1560 1570 1580  
 CTT CTG CCG AGG TGC CAC CGG GCC GAG GCC AAG TAC AGC GGC GTG TGT  
  
 1590 1600 1610 1620 1630  
 25 GCA GGG CGC CGG TGT CGA GGT CGT CGT CGA CGC GCC GGT GTC GAG GTC  
  
 1640  
 GTC GTC GAC-3'  
 30

0238323

wherein A is deoxyadenyl residue; G is a deoxyguanyl residue; C is a deoxycytidyl residue; and T is a deoxythymidyl residue. The structural gene, as indicated above, begins at residue 541 and continues through to residue 1371, terminating with the stop codon located at residue 1372. The amino acid sequence of the tylF structural gene is that indicated under the  
5 corresponding nucleotide sequence. As those skilled in the art will recognize, because of the degeneracy of the genetic code, equivalent sequences to that specifically provided above can be obtained which will encode the same tylF gene product. The means for obtaining such  
10 equivalent sequences will be familiar to those skilled in the art. Further, that a specific sequence is provided is not to be construed as limiting the invention in any way.

Streptomyces fradiae strains can be cultured in a number of ways using any of several different  
15 media. Carbohydrate sources that are preferred in a culture medium include, for example, molasses, glucose, dextran, and glycerol, and nitrogen sources include, for example, soy flour, amino acid mixtures, and peptones. Nutrient inorganic salts are also incorporated into the  
20 medium and include the customary salts capable of yielding sodium, potassium, ammonium, calcium, phosphate, chloride, sulfate, and like ions. As is necessary for the growth and development of other microorganisms, essential trace elements are also added.  
25 Such trace elements are commonly supplied as impurities

incidental to the addition of other constituents of the medium. S. fradiae strains are grown under aerobic culture conditions over a relative wide pH range of about 5.5 to 8 at temperatures ranging from about 25° to 37°C. In particular, tylosin can be produced by cultivation of tylosin-producing strains of, for example, S. fradiae such as those containing the vectors provided by the present invention. The culture medium employed can be any one of a number of media since the organism is capable of utilizing many energy sources. However, for economy of production, maximum yields of antibiotic, and ease of isolation of the antibiotic, certain culture media are preferable. The media which are useful in the production of tylosin include an assimilable source of carbon such as glucose, sucrose, fructose, starch, glycerine, molasses, dextrin, brown sugar, corn steep solids, and the like. The preferred sources of carbon are glucose and starch. Additionally, employable media include a source of assimilable nitrogen such as linseed meal, tankage, fish meal, cotton seed meal, oatmeal, ground wheat, soybean meal, beef extract, peptones (meat or soy), casein, amino acid mixtures, and the like. Preferred sources of nitrogen are soybean meal, casein, and corn steep solids.

Mineral salts, for example, those providing sodium, potassium, ammonium, calcium, magnesium, cobalt, sulfate, chloride, phosphate, carbonate, acetate, and nitrate ions, and a source of growth factors such as distillers' solubles and yeast extract, can be incorporated into the media with beneficial results.

As is necessary for the growth and development of other microorganisms, essential trace elements should also be included in the culture medium for growing the microorganisms employed in this invention. Such trace elements are commonly supplied as impurities incidental to the additional of the other constituents of the medium.

The initial pH of the culture medium can be varied widely. However, it has been found that the initial pH of the medium desirably is between about pH 5.5 and about pH 8.0, and preferably is between about pH 6.5 and about pH 7.0. As has been observed with other organisms, the pH of the medium gradually increases throughout the growth period of the organism during which time tylosin is produced, and may attain a pH from about pH 7.2 to about pH 8.0 or above, the final pH being dependent at least in part on the initial pH of the medium, the buffers present in the medium, and the period of time the organism is permitted to grow.

Submerged, aerobic cultural conditions are the conditions of choice for the production of large amounts of tylosin. For preparation of relatively small amounts, shake flasks and surface culture in bottles can be employed, but for the preparation of large amounts, submerged aerobic culture in sterile tanks is preferred. The medium in the sterile tank can be inoculated with a sporulated suspension. However, because of the growth lag experienced when a sporulated suspension is used in the inoculum, the vegetative form of the culture is preferred to avoid the pronounced growth lag, thereby



permitting a more efficient use of the fermentation equipment. Accordingly, it is desirable first to produce a vegetative inoculum of the organisms by inoculating a relatively small quantity of culture medium with the spore form of the organism, and when a young, active, vegetative inoculum has been obtained, to transfer the vegetative inoculum aseptically to the large tank. The medium in which the vegetative inoculum is produced can be the same or a different medium than that utilized for the large scale production of tylosin.

The organisms grow best at temperatures in a range of about 25°C to about 37°C. Optimal tylosin production appears to occur at a temperature of about 26-30°C.

As is customary in submerged culture processes, sterile air is blown through the culture medium. For efficient growth of the organism and tylosin production, the volume of air employed in the tank production of tylosin preferably is upwards of 0.1 volume of air per minute per volume of culture medium. Efficient growth and optimal yields of tylosin are obtained when the volume of air used is at least one volume of air per minute per volume of culture medium.

The concentration of tylosin activity in the culture medium can readily be followed during the fermentation period by testing samples of the culture medium for their inhibitory activity against the growth of an organism known to be inhibited in the presence of tylosin.

In general, after inoculation, maximum production of tylosin occurs within about 2 to 7 days when submerged aerobic culture or shake flask culture is employed, and within about 5 to 10 days when surface culture is used.

If desired, the mycelium and undissolved solids are removed from the fermentation broth by conventional means such as filtration or centrifugation. If desired, the tylosin is removed from the filtered or centrifuged broth by employing adsorption or extraction techniques familiar to those skilled in the art.

For the extraction of tylosin from the filtered broth, water-immiscible, polar, organic solvents are preferred, such including esters of fatty acids, for example, ethyl acetate and amyl acetate; chlorinated hydrocarbons, for example, chloroform ethylene dichloride, and trichloroethylene; water-immiscible alcohols, for example, butyl and amyl alcohols; water-immiscible ketones, for example, methyl isobutyl ketone and methyl amyl ketone; and others, for example, diethyl ether and methyl propyl ether. Other solvents of similar character can also be employed. Chloroform and amyl acetate are the presently preferred extraction solvents.

For the recovery of tylosin by adsorption techniques, various absorbants and ion exchange resins can be used, for example, carbon, silica gel, alumina, and ion exchange resins of acidic character such as "XE" 64 and "IRC" 50 (weakly acidic cation exchange resins sold by Rohm & Haas Company), carboxymethyl cellulose resin, and "Dowex" 50 (a strongly acidic cation exchange

X-5555

-11-

can be obtained by the Dry Chemical Company. The tylosin can be adsorbed on one of the above or similar adsorbents from a solution in chloroform, acetone, benzene or other suitable solvents. The adsorbed tylosin can then be eluted from the adsorbent by suitable elution techniques such as by washing the adsorbent on which the tylosin is adsorbed with a lower alcohol, for example, methanol or ethanol, or with a lower alcohol containing up to about 50 percent of a lower ketone, for example, acetone.

The organic solvent extract obtained by the preferred extraction method can be directly evaporated to dryness to provide crude tylosin. Alternatively the organic solvent extract can be used to provide purified tylosin by concentration in vacuo the organic solvent extract of tylosin by decolorizing the concentrate with carbon, and by precipitating the tylosin by the addition of a non-polar solvent, for example, petroleum ether. The precipitate which is thus obtained is a solid, purified tylosin which is usually amorphous. The amorphous precipitate can be crystallized by employing one of the crystallizing solvents mentioned above. Alternatively, tylosin can be recovered from a tylosin-containing organic extract, by adsorption chromatography, and by recovery of the adsorbed tylosin from the adsorbent by elution.

Other means for preparing the desired product from the culture medium will be recognized by those skilled in the art.

The acid addition salts of tylosin can be formed with mineral acids, for example, sulfuric, hydrochloride and nitric acid, and with organic acids, for example, tartaric, gluconic, oxalic and acetic acid.

- 5 The acid addition salts can be prepared by dissolving the free base of tylosin in a solvent in which it is soluble, such as acetone or ether, and adding to the solution an equimolar amount of the appropriate acid. The salt which is formed usually precipitates out of  
10 solution. In the event the salt does not precipitate, it can be recovered by evaporating the solution to a smaller volume to permit precipitation, or by adding a miscible solvent in which the salt is not soluble.

- The following non-limited are provided to  
15 further illustrate the invention. Sources of reagents are provided merely for convenience and in no way limit the invention.

#### Example 1

20

#### Isolation of Plasmid pHJL280

##### A. Culture of *E. coli* K12 HB101/pHJL280

- 25 Lyophils of *E. coli* K12 HB101/pHJL280 can be obtained from the NRRL under the accession number NRRL B-18043. The lyophilized cells are streaked onto L-agar plates (L agar contains 10 g of Bacto Tryptone, 5 g of Bacto Yeast Extract, 10 g of NaCl, 2 g of glucose, and

X-6808A

-60-

15 g of agar per liter) containing 50 µg/ml ampicillin to obtain a single-colony isolate of E. coli K12 HB101/pHJL280. One such colony was used to inoculate 100 ml of L broth (L broth is L agar without the agar), which was then incubated aerobically at 37°C overnight (about 16 hours). The following morning, the cells were harvested by centrifugation at 10,000Xg for 10 minutes. The ~1 g of cells obtained by this procedure are used to prepare plasmid pHJL280 DNA in substantial accordance with the procedure described below.

#### B. Plasmid Isolation

The cell pellet obtained in Example 1A was resuspended in 10 ml of a solution composed of 25% sucrose and 50 mM Tris-HCl at a pH = 8.0. About 1 ml of a 10 mg/ml solution of lysozyme in 50 mM Tris-HCl at a pH = 8.0 was added to the cell suspension, and the resulting mixture was incubated on ice for 5 minutes. About 4 ml of 0.25 M EDTA, pH = 8.0, were then added to the cell suspension, and incubation on ice was continued for another 5 minutes. About 16 ml of lysis solution (lysis solution contains 0.4% deoxycholate; 1% Brij58 (Sigma Chemical Co., P.O. Box 14508, St. Louis, MO 63178); 0.05 M Tris-HCl, pH = 8.0; and 0.0625 M EDTA) were added to the lysozyme-treated cells, and the resulting mixture was incubated at 37°C for 15 minutes.

The cell lysate was cleared by centrifugation at 48,000Xg for 25 minutes. The supernatant was decanted into a separate tube, to which was added 0.1 volume

X-6808A

-61-

of 3.0 M NaOAc at a pH = 8.0 and 0.64 volume of isopropyl alcohol. The DNA precipitate was collected by centrifugation at 20,000Xg for 10 minutes and then redissolved in 0.1 volume of TE buffer (10 mM Tris-HCl, pH = 7.8, and 1 mM EDTA). The solution of DNA was incubated at 65°C for 30 minutes and then purified by equilibrium-density-gradient ultracentrifugation in CsCl and propidium diiodide. The plasmid pHJL280 DNA obtained by this procedure was dissolved in TE buffer at a concentration of about 1 µg/µl. A restriction site map of plasmid pHJL280 is presented in Figure 2 of the accompanying drawings.

#### Example 2

15

#### Isolation of Plasmids pHJL284, pHJL309, pHJL311, and pHJL315

Lyophils of the E. coli strains harboring plasmids pHJL284, pHJL309, pHJL311, and pHJL315 can be obtained from the NRRL under the accession numbers listed in Table X. The desired plasmids are each obtained and purified from the lyophilized cells in substantial accordance with the teaching of Example 1. Restriction site maps of the plasmids are presented in Figures 2-6 of the accompanying drawings.

plasmid pHJL280 DNA was added to about 50  $\mu$ l of 1 mg/ml heparin sulfate (Sigma) and incubated on ice for about 10 minutes. Much less plasmid DNA, about 5-100 nanograms, can be used to transform Streptomyces fradiae if prepared from a S. fradiae host. The procedure for isolating Streptomyces plasmid DNA is described in Hopwood et al., 1985, Genetic Manipulation of Streptomyces: A Laboratory Manual (John Innes Foundation, Norwich, England). The DNA/heparin solution was first added to about 200  $\mu$ l of protoplasts, and about 0.9 ml of a solution composed of 55% PEG 1000 (Sigma) in P medium was then added to the DNA/protoplast mixture, and the resulting mixture was gently mixed at room temperature.

The mixture was plated in varying aliquots onto R2 plates using 4 ml of soft-R2-agar overlays. R2 plates contain 30 ml of R2 media and have been dried at 37°C for about 4 days. R2 media is prepared by adding 103 g sucrose, 0.25 g  $K_2SO_4$ , 2 ml of trace element solution, 10.12 g  $MgCl_2 \cdot 6H_2O$ , 10.0 g of glucose, 2.0 g of L-asparagine, 0.1 g of Casamino acids, and 22 g of agar to 700 ml of water; sterilizing the resulting solution; and finally, adding 100 ml of each of the following solutions: 0.05 g  $KH_2PO_4$ /100 ml of deionized water; 2.22 g  $CaCl_2$ /100 ml of deionized water; and 0.25 M TES, pH = 7.2. The pH of the final solution is adjusted to equal 7.2. Trace element solution contains 40 mg  $ZnCl_2$ , 200 mg  $FeCl_3 \cdot 6H_2O$ , 10 mg  $CuCl_2 \cdot 2H_2O$ , 10 mg  $MnCl_2 \cdot 4H_2O$ , 10 mg  $Na_2B_4O_7 \cdot 10H_2O$ , and 10 mg  $(NH_4)_6Mo_7O_{24} \cdot 4H_2O$  per liter. The soft-R2-agar overlays

Example 3Construction of *Streptomyces fradiae* GS28/pHJL280

5           A culture of *Streptomyces fradiae* GS28 was  
inoculated into 20 ml of trypticase-soya broth (TSB) and  
incubated in a water-bath incubator at 29°C at 260 rpm  
overnight (about 16 hours). The culture was homogenized  
using a homogenizing vessel (Thomas Scientific,  
10   Swedesboro, NJ) and a T-Line laboratory stirrer and then  
fragmented using a Sonifier Cell Disruptor (Heat Systems  
Ultrasonics, Inc.) for 7 seconds at 76 Watts. Four ml  
of the homogenized, fragmented culture were inoculated  
into 20 ml of TSB (BBL) containing 0.3% weight by volume  
15   glycine, and the culture was again incubated overnight  
at 29°C. The following morning, the culture was homoge-  
nized and recultured as described above. After this  
third overnight incubation, the culture was homogenized,  
collected, and then washed twice with P media. P media  
20   was prepared by adding 103 g of sucrose to 0.25 g of  
 $K_2SO_4$  and 2.03 g of  $MgCl_2 \cdot 6H_2O$  and then adding deionized  
water to a final volume of 700 ml. The mixture was then  
sterilized, and to each 70 ml of solution, about 10 ml  
each of 0.05 g  $KH_2PO_4$ /100 ml of deionized water; 2.78 g  
25    $CaCl_2$ /100 ml of deionized water; and 0.25 M TES  
(2-([tris-(hydroxymethyl)methyl]-amino)ethanesulfonic  
acid)) at a pH = 7.2 were added.

The cell pellet was resuspended in 15 ml of P  
media containing 1 mg/ml lysozyme (Calbiochem, La Jolla,  
30   CA 92037) and then incubated at room temperature for  
about one-and-one-half hours to form protoplasts. The  
protoplasts were gently collected by centrifugation,  
washed twice with P media, r suspend d in 2 ml of P  
media, and incubated on ice until use. About 1 µg of



are prepared by adding 51.5 g of sucrose, 5.06 g  $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ , 1.11 g  $\text{CaCl}_2$ , 50 ml of 0.25 M TES at a pH = 7.2, and 2.05 g agar to enough deionized water to achieve a final volume of 500 ml. The mixture is steamed to melt the agar, decanted into 4 ml aliquots, and autoclaved prior to use. After the transformed protoplasts had been plated, the plates were incubated at 29°C for 24 hours, and then, 4 ml of soft-R2 agar containing 25  $\mu\text{l}$  of 50 mg/ml thiostrepton (E. R. Squibb, Princeton, NJ 08540) were spread over the protoplasts. Incubation of the plates at 29°C was continued until regeneration was complete, usually a period of about 7-14 days, to select for the desired S. fradiae GS28/pHJL280 transformants.

The Streptomyces fradiae GS28/pHJL280 strain was cultured and produced macrocin O-methyltransferase and tylosin at levels above that produced in the untransformed S. fradiae GS28 strain. Macrocin O-methyltransferase activity was assayed and determined in substantial accordance with the teaching of Yeh et al., 1984, Journal of Chromatography 288:157-165. Comparison of the macrocin O-methyltransferase activities in the transformed, GS28/pHJL280, and parental, GS28, strains showed a 60-to-100-fold increase of enzyme and 14-to-18-fold increase of tylosin production in the transformed strain. Tylosin production was assayed and determined in substantial accordance with the teaching of Baltz and Seno, 1981, Antimicrobial Agents and Chemotherapy 20:214-225; and Kennedy, J.H., 1983, Journal of Chromatography 281:288-292.

Example 4Construction of *Streptomyces fradiae* GS15/pHJL280

5           The desired strain was constructed in substantial accordance with the teaching of Example 3 except that *Streptomyces fradiae* GS15, rather than *S. fradiae* GS28, was used. The desired strain was cultured for 72 hours and produced macrocin Q-methyltransferase and  
10   tylosin at levels above that produced in the untransformed *S. fradiae* GS15 strain, which produces no readily detectable tylosin.

Example 5

15

Construction of *Streptomyces fradiae* GS15/pHJL284

          The desired strain was constructed in substantial accordance with the teaching of Example 4 except  
20   that plasmid pHJL284, rather than plasmid pHJL280, was used. The desired strain was cultured and produced macrocin Q-methyltransferase and tylosin at levels above that produced in the untransformed *S. fradiae* GS15  
strain.

25

0238323

-66-

0238323

Example 6Construction of *Streptomyces fradiae* GS16/pHJL280

The desired strain was constructed in substantial accordance with the teaching of Example 3 except *Streptomyces fradiae* GS16, rather than *S. fradiae* was used. The desired strain was cultured and produced the tylE gene product, demethylmacrocin methyltransferase, and tylosin at levels above that produced in the untransformed strain. The demethylmacrocin O-methyltransferase activity and tylosin production respectively are assayed and determined in substantial accordance with the well-known referenced procedures, except that demethylmacrocin is substituted for macrocin as substrate.

Example 7Construction of *Streptomyces fradiae* GS76/pHJL280

The desired strain was constructed in substantial accordance with the teaching of Example 3 except *Streptomyces fradiae* GS76, rather than *S. fradiae* was used. The desired strain was cultured and produced the tylD and tylH gene products and tylosin at levels above that produced in the untransformed strain.

L280

substan-  
cept  
*fradiae*  
and  
els

L284

substan-  
cept  
4,  
were  
ed the  
at

and  
Method

ffec-  
ts  
l  
results  
rom  
d Seno,

2285

87265660

- 1981, Antimicrobial Agents and Chemotherapy 20:214-225)  
that also contained 20 µg/ml thiostrepton if the strain  
being cultured harbored a plasmid. Note that the  
transformed strains listed in Tables XII and XIII are  
5 low tylosin-producing, or produce amounts of tylosin  
that are not readily detectable, and were cultured in  
the presence of selective pressure (thiostrepton) for  
plasmid maintenance as an autonomously replicating  
vector.

0238323

**Table XII**  
**Specific Activity of the tylF Gene Product, Macrocin O-methyltransferase (MOMT)**

|       | Strain             | Transforming Plasmid | MOMT Specific Activity |        |        |        |
|-------|--------------------|----------------------|------------------------|--------|--------|--------|
|       |                    |                      | 2 days <sup>1</sup>    | 3 days | 4 days | 6 days |
| Run 1 | GS15               | pHJL210 <sup>4</sup> | 0                      | 0      | 0      | 0      |
|       | GS15               | pHJL280              | 1.14                   | 1.93   | NT     | NT     |
|       | C4 <sup>2</sup>    | None                 | Not tested (NT)        | 0.35   | 0.16   | NT     |
| Run 2 | GS15               | pHJL210              | 0                      | 0      | 0      | 0      |
|       | GS15               | pHJL280              | 4.2                    | 3.2    | 2.2    | 1.8    |
|       | C4                 | None                 | 0.8                    | 1.0    | 0.9    | 0.9    |
|       | T1405 <sup>3</sup> | None                 | 0.9                    | 1.2    | 1.5    | 1.4    |
| Run 3 | GS28               | None                 | 0                      | 0.01   | 0.03   | NT     |
|       | GS28               | pHJL210              | 0                      | 0      | 0      | NT     |
|       | GS28               | pHJL280              | 0.8                    | 0.7    | 1.0    | NT     |
|       | GS28               | pHJL284              | 0.9                    | 1.2    | 0.9    | NT     |
|       | C4                 | None                 | 0.2                    | 0.6    | 0.5    | NT     |

<sup>1</sup>days in fermentation.<sup>2</sup>the strain from which GS15 and GS28 were derived.<sup>3</sup>a strain derived from C4.<sup>4</sup>the cloning vector into which the tyl genes were inserted to obtain plasmids pHJL280 and pHJL284.

2288

87265660

Table XIII

Specific Activity of the tyle Gene Product, Demethylmacrocin  
O-methyltransferase (DMOMT)

| Strain | Transforming<br>Plasmid | DMOMT Specific Activity |        |        |
|--------|-------------------------|-------------------------|--------|--------|
|        |                         | 2 days*                 | 3 days | 4 days |
| GS16   | pHJL210                 | 0                       | 0      | 0      |
| GS16   | pHJL280                 | 1.8                     | 3.7    | 4.0    |
| GS16   | pHJL280                 | 3.8                     | 1.7    | 3.0    |
| GS16   | pHJL284                 | 1.3                     | 1.6    | 2.2    |
| C4     | pHJL210                 | 0.7                     | 1.3    | 1.5    |
| C4     | pHJL210                 | 0.2                     | 1.1    | 1.9    |
| C4     | None                    | 0.4                     | 1.5    | 1.0    |

The results in Table XIV were obtained from transformants of high tylosin-producing strains that were cultured post-transformation to obtain integrants, transformants in which all or part of the plasmid DNA has integrated into the genome of the host cell. Two methods were used to obtain the integrants. In the first method, transformants are passaged onto selective (contains thiostrepton) and nonselective plates and incubated about 16 hours at 29°C to obtain single colonies. The single colonies on the nonselective plates that were thiostrepton-resistant on the selective plate are repassaged several times in the same manner until a single colony was found to be relatively stable without selection. In the second method for obtaining integrants, the transformants were nonselectively passaged several times by transferring spores from the surface of the plate using a cotton swab. After several passages, the colonies are grown in non-selective, liquid media (TSB), homogenized, fragmented by sonication, diluted, and plated on selective and nonselective media to identify relatively stable integrants. Other methods of obtaining integrants are apparent to those skilled in the art, and the present method is not limited to a particular method of obtaining integrants.

Relatively stable integrants were used to inoculate vegetative medium (complex vegetative medium contains, per liter, 10 g of corn steep liquor, 5 g of yeast extract, 5 g of soybean grits, 3 g of calcium carbonate, and 4.5 g of crude soybean oil, and the pH is adjusted to 7.8 with NaOH. TSB is also a suitable

vegetative media) without thiostrepton (no selective pressure), and the vegetative culture was used to inoculate (10% inoculum) the fermentation medium, which also lacked thiostrepton. Fermentations were run at 260 rpm at 29°C for seven days. The total macrolide content of the fermentation broth was measured by extraction with methanol:CHCl<sub>3</sub>, reading the absorbance at 290 nm, and comparing to a standard curve. Tylosin factors were identified by spotting the fermentation broth onto silica-gel-TLC plates and developing the plates with a solvent system of 95:5 ethylacetate:diethylamine. The concentration of individual macrolide components was the total A<sub>290</sub> times the percentage of each component as determined by HPLC.

15



Table XIV

| Strain | Thiostrepton Resistant | Transforming Plasmid | % DMOMT Specific Activity |      | MOMT Specific Activity | Tylosin* |
|--------|------------------------|----------------------|---------------------------|------|------------------------|----------|
|        |                        |                      | DMOMT                     | %    |                        |          |
| C4     | 0                      | None                 | 0.59                      | 0.14 | 1                      | 1        |
| C4     | 9.5                    | pHJL280              | Not tested (NT)           | NT   | 1.10                   | 1.10     |
| C4     | 5.9                    | pHJL280              | NT                        | NT   | 0.97                   | 0.97     |
| T1405  | 0                      | None                 | 1.0                       | 0.17 | 1.14                   | 1.14     |
| T1405  | 50                     | pHJL280              | 0.91                      | 0.26 | 1.52                   | 1.52     |
| T1405  | 8.7                    | pHJL280              | NT                        | NT   | 1.21                   | 1.21     |
| T1405  | 11                     | pHJL280              | NT                        | NT   | 1.07                   | 1.07     |
| T1405  | 6.4                    | pHJL280              | NT                        | NT   | 1.21                   | 1.21     |
| T1405  | 18                     | pHJL280              | 2.5                       | 0.43 | 1.60                   | 1.60     |
| T1405  | 4.6                    | pHJL280              | NT                        | NT   | 0.98                   | 0.98     |
| T1405  | 16                     | pHJL280              | NT                        | NT   | 1.07                   | 1.07     |
| T1405  | 12                     | pHJL280              | NT                        | NT   | 1.10                   | 1.10     |
| T1405  | 18                     | pHJL280              | NT                        | NT   | 1.28                   | 1.28     |
| T1405  | 25                     | pHJL280              | NT                        | NT   | 1.28                   | 1.28     |
| T1405  | 56                     | pHJL280              | 0.82                      | 0.22 | 1.45                   | 1.45     |

\*Relative to C4 strain

Example 11Preparation of Tylosin

5           A sporulated culture of *Streptomyces fradiae*  
containing the plasmids provided by the present inven-  
tion can be produced by growing the organism on a  
nutrient agar slant having the following composition:

|    |  |       |
|----|--|-------|
| 10 | Yeast extract  | 1.0 g |
|    | Beef extract   | 1.0 g |
|    | Hydrolyzed casein ("N-Z-Amine-Type<br>A," sold by the Sheffield Chemical<br>Co.) | 2 g   |
| 15 | Dextrin  | 10 g  |
|    | Cobaltous chloride heptahydrate  | 20 mg |
|    | Agar   | 20 g  |
|    | Water  | 1 l.  |

The pH of the medium is adjusted to pH 7.3 by the addition of sodium hydroxide.

The slant is inoculated with spores of the desired organism and is incubated for five days at about 30°C. The sporulated culture growth on the slant is covered with water, and the slant is scraped gently to remove the spores to provide an aqueous spore suspension.

1 ml. of the spore suspension is used to inoculate under aseptic conditions a 100 ml. portion of a sterile vegetative culture medium having the following composition:

|    |  |          |
|----|--|----------|
| 15 | Glucose  | G.<br>15 |
|    | Soybean meal                                       | 15       |
|    | Corn steep solids                                  | 15       |
|    | Sodium chloride                                    | 5        |
|    | Calcium carbonate                                  | 2        |
| 20 | Tap water, added to make a total of volume of 1 l. |          |

The inoculated vegetative medium is incubated at about 30°C. for 48 hours, during which time the incubate is shaken at the rate of 114 cycles per minute on a reciprocal shaker having a 2-inch stroke.

5 ml. of the vegetative inoculum are used to inoculate aseptically 100 ml. portions of the following sterilized production medium contained in 500 ml. Erlenmeyer flasks:

|    |  |       |
|----|--|-------|
| 35 | Soybean meal                                       | 15 g  |
|    | Casein   | 1 g   |
|    | Crude glucose syrup                                | 20 ml |
|    | Calcium carbonate                                  | 2.5 g |
|    | Sodium nitrate                                     | 3 g   |
|    | Tap water, added to make a total of volume of 1 l. |       |

The inoculated culture then is incubated for 100 hours at about 26-28°C. During the incubation period, the incubate is shaken at 114 revolutions per minute on a reciprocal shaker having a 2-inch stroke.

5 The pH of the starting medium is about pH 6.5, and at the end of the incubation period, the pH of the medium generally increases to about pH 7.5.

The fermented culture broth is filtered to remove the mycelium and other undissolved solids. The

10 filtered broth contains the tylosin.

#### Example 12

#### Alternate Preparation of Tylosin

15

A sporulated culture of the desired transformed microorganism is produced by growing the organism on a nutrient agar slant having the following composition.

20

|   |    |
|---|----|
| Tomato paste-oatmeal Agar:                  | G  |
| Tomato paste                                | 20 |
| Pre-cooked oatmeal                          | 20 |
| Agar  | 15 |
| Water, added to make a total volume of 1 l. |    |

25

The slant is inoculated with spores of the organism and the inoculated slant is incubated for 9

days at a temperature of about 30°C. After incubation, the sporulated culture on the slant is covered with water, and the surface of the slant is scraped gently to remove the spores to obtain an aqueous spore suspension.

5           Employing aseptic techniques, one-half of the inoculum obtained from one agar slant is used for inoculating a 500 ml. portion of a sterilized vegetative culture medium having the following composition contained in a 2 l. Erlenmeyer flask:

10

## Corn-steep yeast I:

|   |    |
|---|----|
| Glucose                                     | G. |
| Corn steep solids                           | 15 |
| Yeast                                       | 5  |
| Calcium carbonate                           | 5  |
| Water, added to make a total volume of 1 l. | 5  |

15

20           The incubation is carried on at 28°C for 48 hours with shaking at 110 cycles per minute on a reciprocal shaker having a 2-inch stroke.

0.25 gal. of the vegetative inoculum from the flask is added aseptically as an inoculum to 250 gal. of the sterile corn steep yeast I medium described above contained in an iron 350 gal fermentor. 0.025 gal. of Antifoam A (an antifoam product sold by The Dow Corning Company) is added to the culture medium to prevent excessive foaming, and additional quantities are added as needed during the fermentation. The inoculated medium is fermented for 24 hours at a temperature of 28°C. During fermentation, the medium is aerated with sterile air at a rate of 27 cubic feet per minute and is

25

30

agitated with two 16-inch impellers operated at 160 revolutions per minute.

To a 1700 gal. iron fermentor are added 1200 gal. of a medium having the following composition.

|    |  |    |     |
|----|--|----|-----|
| 5  |  |    |     |
|    | Corn steep soy XII:                            |    | Kg. |
|    | • Glucose                                      | 30 |     |
|    | • Soybean oil meal                             | 15 |     |
|    | Corn steep solids                              | 5  |     |
| 10 | Crude soybean oil                              | 10 |     |
|    | Calcium carbonate                              | 2  |     |
|    | Sodium chloride                                | 5  |     |
|    | Water, added to make a total volume of 1000 l. |    |     |

15 The medium is inoculated with 96 gal. of the inoculum grown in the fermentation tank. The fermentation is carried on at 28°C for four days, and any foaming is controlled by the addition as needed of "Larex" No. 1 (an antifoam product sold by Swift & Company). The fermentation medium is aerated by the addition of sterile air at the rate of 128 cubic feet per minute and is agitated with two 24-inch impellers operated at 130 revolutions per minute.

25 600 pounds of "Silflo" (a diatomaceous earth filter aid sold by The Silfo Company) are added to the broth, and the mixture is filtered. The filtrate is adjusted to pH 8.5 by the addition of 20 percent sodium hydroxide, 500 gal. of chloroform are added, the mixture is stirred for 30 minutes, and the chloroform layer which is in the form of an emulsion is decanted. The chloroform extraction is repeated twice with 500-gal. portions of chloroform. The chloroform emulsions which

contain the tylosin are combined and are passed through a De Laval separator to break the emulsion, and the chloroform solution is then concentrated in vacuo to a volume of 25 l. The impurities are largely removed from the solution by passing it over a column 6 inches in diameter containing 10 kg. of activated carbon such as that sold by the Pittsburgh Coke and Chemical Co. The carbon column is washed with 16 l. of chloroform, and the combined chloroform effluents containing the tylosin are concentrated in vacuo to a volume of about 2 l. The chloroform concentrate is added slowly with stirring to 20 l. of petroleum ether, the mixture is stirred for 15 minutes, it is filtered to remove the white, amorphous precipitate of tylosin.

The amorphous tylosin is crystallized by dissolving it in 355 ml. of acetone, filtering the acetone mixture to remove a slight haze, and slowly adding the filtered acetone mixture with gentle stirring to 20 l. of water at 5°C. The aqueous, acetone solution of tylosin is permitted to stand at room temperature with gentle stirring to permit the acetone to evaporate slowly, whereupon tylosin crystallizes out. The tylosin crystals are removed by filtration and are dried in vacuo at room temperature. Tylosin has a melting point of about 127-132°C.

Example 13Preparation of Tylosin Tartrate

5           5 g. of crystalline tylosin are dissolved in  
100 ml. of acetone, and 1.5 g. of D-tartaric acid  
dissolved in 20 ml. of acetone are added with stirring.  
The solution is permitted to stand at room temperature  
whereupon the tartrate salt of tylosin crystallizes out  
10 of the solution. The crystals of the tartrate salt of  
tylosin are removed by filtration, are washed with  
acetone, and are air-dried. The crystalline tartrate  
salt of tylosin melts at about 140-146°C.

Example 14Preparation of Tylosin Gluconate

15           1.03 g. glucono-delta lactone are dissolved in  
20 10 ml. of water, and the aqueous solution is warmed to  
85°C. for two hours to cause hydrolysis of the lactone  
to gluconic acid. 15 ml. of warm methanol are added to  
the aqueous solution. 5 g. of tylosin dissolved in 10  
ml. of methanol are added to the methanol mixture with  
25 stirring. The tylosin methanol mixture is permitted to  
stand overnight at room temperature. The methanol is  
removed from the mixture by evaporation in vacuo at room  
temperature. After the methanol is removed, 40 ml. of  
water are added to the aqueous tylosin mixture. The  
30 diluted mixture is filtered, and the filtrate containing  
the tylosin is freeze-dried, producing a white solid  
consisting of the gluconate salt of tylosin. Tylosin  
gluconate salt melts at about 114-117°C.



Example 15Preparation of Tylosin Hydrochloride

5           890 mg. of tylosin are dissolved in 200 ml. of  
ether. The ether mixture is acidified by the addition  
of 0.082 ml. of 12 N hydrochloric acid. The precipitate  
of the hydrochloride salt of tylosin which forms is  
filtered off, is washed with ether, and is dried in  
10 vacuo. The hydrochloride salt of tylosin is  
recrystallized from an ethanol-ether mixture. The  
hydrochloride salt of tylosin has melting point of about  
141-145°C.

Claims

1. A method for increasing the antibiotic- or  
antibiotic precursor-producing ability of an  
5 antibiotic-producing microorganism, which comprises  
culturing a microorganism which produces an antibiotic  
or antibiotic precursor by a biosynthetic pathway, said  
microorganism being transformed with a DNA cloning  
10 vector or portion thereof which contains an antibiotic  
or antibiotic-precursor biosynthetic gene coding for  
expression of a rate-limiting enzyme or gene product of  
the biosynthetic pathway, under conditions suitable for  
cell growth, expression of the antibiotic or  
15 antibiotic-precursor biosynthetic gene and production of  
the antibiotic or antibiotic precursor, provided that  
the culturing process provides an increase in the  
antibiotic-producing ability of the microorganism.

2. A method as claimed in Claim 1 in which  
the microorganism is Streptomyces, Cephalosporium, or  
20 Penicillium.

3. A method as claimed in Claim 1 or 2 in  
which the microorganism, antibiotic, antibiotic precursor,  
antibiotic biosynthetic pathway, and antibiotic  
biosynthetic gene respectively are Streptomyces, tylo-  
25 sin, a tylosin precursor, tylosin biosynthetic pathway,  
and a tylosin biosynthetic gene.

4. A method as claimed in Claim 3 in which  
the microorganism is S. fradiae, S. rimosus, and S.  
hygroscopicus.

5. A method as claimed in Claim 4 in which the tylosin biosynthetic gene is tylC, tylD, tylE, tylF, tylH, tylJ, tylK, tylL, or tylM.

6. A method as claimed in any one of Claims 1 to 5 in which the biosynthetic gene is tylF.

7. A method as claimed in Claim 1 in which the cloning vector is plasmid pHJL280, pHJL284, pHJL309, pHJL311, or pHJL315.

8. A method as claimed in claim 1 in which the microorganism transformed with a cloning vector is *Streptomyces fradiae* GS15/pHJL280, *S. fradiae* GS15/pHJL284, *S. fradiae* GS15/pHJL311, *S. fradiae* GS15/pHJL315, *S. fradiae* GS28/pHJL280, *S. fradiae* GS28/pHJL284, *S. fradiae* GS28/pHJL311, *S. fradiae* GS28/pHJL315, *S. fradiae* GS16/pHJL280, *S. fradiae* GS16/pHJL315, *S. fradiae* GS48/pHJL280, *S. fradiae* GS48/pHJL315, *S. fradiae* GS52/pHJL284, *S. fradiae* GS52/pHJL311, *S. fradiae* GS76/pHJL280, or *S. fradiae* GS76/pHJL315.

9. A recombinant DNA cloning vector as defined in claim 1.

10. Plasmids pHJL280, pHJL284, pHJL309, pHJL311, or pHJL315.

11. A microorganism that is transformed with a cloning vector as claimed in claim 9.

12. A microorganism which is transformed with a plasmid of Claim 10.

13. A microorganism of Claim 12 which is *E. coli* K12 HB101/pHJL280, *E. coli* K12 HB101/pHJL284, *E. coli* K12 HB101/pHJL309, *E. coli* K12 HB101/pHJL311, or *E. coli* K12 JM109/pHJL315.

14. A microorganism as claimed in Claim 12 which is *Streptomyces*.

15. A microorganism as claimed in Claim 14 which is *Streptomyces fradiae*.

5 16. *S. fradiae* GS15/pHJL280, *S. fradiae* GS15/pHJL284, *S. fradiae* GS15/pHJL311, *S. fradiae* GS15/pHJL315, *S. fradiae* GS28/pHJL280, *S. fradiae* GS28/pHJL284, *S. fradiae* GS28/pHJL311, *S. fradiae* GS28/pHJL315, *S. fradiae* GS16/pHJL315, *S. fradiae* GS48/pHJL280, *S. fradiae* GS48/pHJL315, *S. fradiae* GS52/pHJL284, *S. fradiae* GS52/pHJL311, *S. fradiae* GS76/pHJL280, or *S. fradiae* GS76/pHJL315.

17. A DNA sequence that encodes a gene for the tylC, tylD, tylE, tylF, tylH, tylJ, tylK, tylL, or tylM biosynthetic genes.

18. A DNA sequence which encodes the promoter and translational-activating sequence of the tylC, tylD, tylE, tylF, tylH, tylJ, tylK, tylL, or tylM biosynthetic genes.

20 19. A DNA sequence which encodes the promoter and translational-activating sequence of the tylF gene.

20. A DNA sequence as claimed in Claim 19 encoding the tylF promoter that is

5'-CTG TGT CAG GTC GCC CGT GGT GAC GGG CTC CGG GGC GGC GGC GCG

25

R

GGC GGC CGA CCT TGA CAT ACC CGC GGC CGG GCT CCT CGT TCC GGC GCG

GCC CGC GCC GAT AGC GTC CGT CCT CAC CGG CTC CGG CGT CCG CGT CCC

CGC CGG GAC GTG CCA CCT CTC CCG ACC CCG CGA GCC GAT CGA CCC GCT

30

ACT GGA GGA CCC-3'

wherein A is a deoxyadenyl residue; G is a deoxyguanyl residue; C is a deoxycytidyl residue; and T is a thymidyl residue and wherein R is a sequence of deoxyribonucleotide that is complementary to said DNA sequence depicted such that A is paired with T; T is paired with A; G is paired with C; and C is paired with G.

21. A process for preparing an antibiotic, an antibiotic precursor, or a pharmaceutically acceptable salt thereof, which comprises culturing a microorganism which produces an antibiotic or antibiotic precursor through an antibiotic biosynthetic pathway, said microorganism being transformed with a DNA cloning vector, or portion thereof, in a culture medium containing assimilable sources of carbon, nitrogen and inorganic salts under aerobic fermentation conditions characterized in that the DNA cloning vector, or portion thereof, comprises an antibiotic biosynthetic gene which codes for the expression of a rate-limiting enzyme or gene product of the antibiotic biosynthetic pathway, said antibiotic biosynthetic gene being expressed under fermentation conditions providing for an increase in the antibiotic-producing ability of the microorganism.

22. A process as claimed in Claim 21 in which the antibiotic or antibiotic-precursor is tylosin or a tylosin-precursor.

5

23. A DNA sequence encoding the tylF gene.

24. The tylF gene.

25. The gene claimed in claim 24 having the sequence:

10                   10                   20                   30                   40  
5'-TTC GCG GGA TGG ATG CTG ACC CGG GGG TCG GCC AGC AGC GCC CGG ACG

                  50                   60                   70                   80                   90  
TGA TCT GGC GGG AGA TCA GCC AGA CCG GCG CCC CGT CCC ACA GCT CGG

15                   100                   110                   120                   130                   140  
CCC GGG CGA TCG GCT CCT CCG CCC GGA GGG CGG CGT ACT GCT CGG GAG

                  150                   160                   170                   180                   190  
GGC TGA AGG GAC AGG TGC GGG CGA CCG GCC AGG CGA TGC TGC GCC GGC

20                   200                   210                   220                   230                   240  
CTG CGG CCC CGT CGG TGT CGT TGG CGC GTG CTG CGG GCA ACA GAA TCC

                  250                   260                   270                   280  
CCT TTT GTG ACG GGC GGG CGT CCC CGG ACG AGG ACA CGA CTC GCT GCG

                  290                   300                   310                   320                   330  
GCC TCA ACG AAA ACA CCG TGT CCG GTG CCC AGG CCA CGA ACG GTG ACC

30                   340                   350                   360                   370                   380  
GGT CTG TGT CAG GTC GCC CGT GGT GAC GGG CTC CGG GGC GGC GGC GCG

wherein A is a deoxyadenyl residue; G is a deoxyguanyl residue; C is a deoxycytidyl residue; and T is a thymidyl residue and wherein R is a sequence of deoxyribonucleotide that is complementary to said DNA sequence depicted such that A is paired with T; T is paired with A; G is paired with C; and C is paired with G.

21. A process for preparing an antibiotic, an antibiotic precursor, or a pharmaceutically acceptable salt thereof, which comprises culturing a microorganism which produces an antibiotic or antibiotic precursor through an antibiotic biosynthetic pathway, said microorganism being transformed with a DNA cloning vector, or portion thereof, in a culture medium containing assimilable sources of carbon, nitrogen and inorganic salts under aerobic fermentation conditions characterized in that the DNA cloning vector, or portion thereof, comprises an antibiotic biosynthetic gene which codes for the expression of a rate-limiting enzyme or gene product of the antibiotic biosynthetic pathway, said antibiotic biosynthetic gene being expressed under fermentation conditions providing for an increase in the antibiotic-producing ability of the microorganism.

22. A process as claimed in Claim 21 in which the antibiotic or antibiotic-precursor is tylosin or a tylosin-precursor.

5

23. A DNA sequence encoding the tylF gene.

24. The tylF gene.

25. The gene claimed in claim 24 having the sequence:

10                   10                   20                   30                   40  
5'-TTC GCG GGA TGG ATG CTG ACC CGG GGG TCG GCC AGC AGC GCC CGG ACG

                  50                   60                   70                   80                   90  
TGA TCT GGC GGG AGA TCA GCC AGA CCG GCG CCC CGT CCC ACA GCT CGG

15                   100                   110                   120                   130                   140  
CCC GGG CGA TCG GCT CCT CCG CCC GGA GGG CGG CGT ACT GCT CGG GAG

                  150                   160                   170                   180                   190  
GGC TGA AGG GAC AGG TGC GGG CGA CCG GCC AGG CGA TGC TGC GCC GGC

20                   200                   210                   220                   230                   240  
CTG CGG CCC CGT CGG TGT CGT TGG CGC GTG CTG CGG GCA ACA GAA TCC

                  250                   260                   270                   280  
CCT TTT GTG ACG GGC GGG CGT CCC CGG ACG AGG ACA CGA CTC GCT GCG

25                   290                   300                   310                   320                   330  
GCC TCA ACG AAA ACA CCG TGT CCG GTG CCC AGG CCA CGA ACG GTG ACC

30                   340                   350                   360                   370                   380  
GGT CTG TGT CAG GTC GCC CGT GGT GAC GGG CTC CGG GGC GGC GGC GCG



0238323

390 400 410 420 430  
GGC GGC CGA CCT TGA CAT ACC CGC GGC CGG GCT CCT CGT TCC GGC GCG

5 440 450 460 470 480  
GCC CGC GCC GAT AGC GTC CGT CCT CAC CGG CTC CGG CGT CCG CGT CCC

490 500 510 520  
CGC CGG GAC GTG CCA CCT CTC CCG ACC CCG CGA GCC GAT CGA CCC GCT

10 530 540 550 560 570  
ACT GGA GGA CCC GTG GCA CCT TCC CCG GAC CAC GCC CGC GAT CTC TAC  
VAL ALA PRO SER PRO ASP HIS ALA ARG ASP LEU TYR  
5 10

15 580 590 600 610 620  
ATC GAG CTG CTG AAG AAG GTC GTC TCG AAC GTC ATC TAC GAG GAC CCC  
ILE GLU LEU LEU LYS LYS VAL VAL SER ASN VAL ILE TYR GLU ASP PRO  
15 20 25

20 630 640 650 660 670  
ACC CAT GTG GCG GGG ATG ATC ACC GAC GCG TCG TTC GAC CGG ACG TCC  
THR HIS VAL ALA GLY MET ILE THR ASP ALA SER PHE ASP ARG THR SER  
30 35 40

25 680 690 700 710 720  
CGT GAG AGC GGC GAG GAC TAC CCC ACG GTC GCC CAC ACG ATG ATC GGC  
ARG GLU SER GLY GLU ASP TYR PRO THR VAL ALA HIS THR MET ILE GLY  
45 50 55 60

2306

87265660

0238323

|    |   |     |     |      |
|----|---|-----|-----|------|
|    | 730   | 740 | 750 | 760  |
|    | CTC AAG CGT CTG GAC AAT CTC CAC CGG TGC CTC GCG GAC GTC GTG GAG |     |     |      |
|    | LEU LYS ARG LEU ASP ASN LEU HIS ARG CYS LEU ALA ASP VAL VAL GLU |     |     |      |
| 5  | 65  |     | 70  | 75   |
|    | 770   | 780 | 790 | 800  |
|    | GAC GGC GTC CCC GGT GAC TTC ATC GAG ACC GGG GTG TGC CGC GCG CCG |     |     |      |
|    | ASP GLY VAL PRO GLY ASP PHE ILE GLU THR GLY VAL CYS ARG ALA PRO |     |     |      |
| 10 | 80  |     | 85  | 90   |
|    | 820   | 830 | 840 | 850  |
|    | TGC ATC TTC GCC CGC GGA CTG CTG AAC GCG TAC GGC CAG GCC GAC CGC |     |     |      |
|    | CYS ILE PHE ALA ARG GLY LEU LEU ASN ALA TYR GLY GLN ALA ASP ARG |     |     |      |
| 15 | 95  |     | 100 | 105  |
|    | 870   | 880 | 890 | 900  |
|    | ACC GTC TGG GTC GCC GAC TCC TTC CAG GGC TTT CCC GAG CTG ACC GGG |     |     |      |
|    | THR VAL TRP VAL ALA ASP SER PHE GLN GLY PHE PRO GLU LEU THR GLY |     |     |      |
| 20 | 110   |     | 115 | 120  |
|    | 920   | 930 | 940 | 950  |
|    | TCC GAC CAC CCG CTG GAC GTC GAG ATC GAC CTC CAC CAG TAC AAC GAG |     |     |      |
|    | SER ASP HIS PRO LEU ASP VAL GLU ILE ASP LEU HIS GLN TYR ASN GLU |     |     |      |
| 25 | 125   |     | 130 | 135  |
|    | 970   | 980 | 990 | 1000 |
|    | GCC GTG GAC CTG CCC ACC AGC GAG GAG ACC GTG CCG GAG AAC TTC GCC |     |     |      |
|    | ALA VAL ASP LEU PRO THR SER GLU GLU THR VAL ARG GLU ASN PHE ALA |     |     |      |
| 30 | 145   |     | 150 | 155  |

2307

87265660

0238323

|    |   |      |      |      |      |
|----|---|------|------|------|------|
|    | 1010  | 1020 | 1030 | 1040 | 1050 |
|    | CGG TAC GGG CTG CTC GAC GAC AAC GTC CGT TTC CTG GCG GGG TGG TTC |      |      |      |      |
|    | ARG TYR GLY LEU LEU ASP ASP ASN VAL ARG PHE LEU ALA GLY TRP PHE |      |      |      |      |
| 5  |   | 160  | 165  |      | 170  |
|    | • 1060  | 1070 | 1080 | 1090 | 1100 |
|    | AAG GAC ACC ATG CCG GCT GCG CCC GTG AAG CAG CTC GCG GTG ATG CGC |      |      |      |      |
|    | LYS ASP THR MET PRO ALA ALA PRO VAL LYS GLN LEU ALA VAL MET ARG |      |      |      |      |
| 10 |   | 175  | 180  |      | 185  |
|    | 1110  | 1120 | 1130 | 1140 | 1150 |
|    | CTG GAC GGC GAC TCC TAC GGC GCC ACC ATG GAT GTG CTC GAC AGC CTG |      |      |      |      |
|    | LEU ASP GLY ASP SER TYR GLY ALA THR MET ASP VAL LEU ASP SER LEU |      |      |      |      |
|    | 190   | 195  |      | 200  |      |
|    | 1160  | 1170 | 1180 | 1190 | 1200 |
|    | TAC GAG CGG CTG TCG CCG GGC GGT TAC GTC ATC GTC GAC GAC TAC TGC |      |      |      |      |
|    | TYR GLU ARG LEU SER PRO GLY GLY TYR VAL ILE VAL ASP ASP TYR CYS |      |      |      |      |
| 20 | 205   | 210  | 215  |      | 220  |
|    | 1210  | 1220 | 1230 | 1240 |      |
|    | ATC CCG GCC TGC CGC GAG CCG TGC ACG ACT TCC GCG ACC GGC TCG GCA |      |      |      |      |
|    | ILE PRO ALA CYS ARG GLU ARG CYS THR THR SER ALA THR GLY SER ALA |      |      |      |      |
| 2  |   | 225  | 230  |      | 235  |
|    | 1250  | 1260 | 1270 | 1280 | 1290 |
|    | TCC GCG ACA CGA TCC ACC GGA TCG ACC GCC AGG GCG CTA TTG GCG GCA |      |      |      |      |
|    | SER ALA THR ARG SER THR GLY SER THR ALA ARG ALA LEU LEU ALA ALA |      |      |      |      |
| 30 |   | 240  | 245  |      | 250  |

0238323

5                   1300                   1310                   1320                   1330                   1340  
CAG CGG CTG AGT CGT TCC GCC CGA GAG CCC GAC GAG AGC AGG AGA TAT  
GLN ARG LEU SER ARG SER ALA ARG GLU PRO ASP GLU SER ARG ARG TYR  
                  255                   260                   265

10                   1350                   1360                   1370                   1380                   1390  
GCG AGA CAC GAC GCG CCC GCT CGG CAT TGA GGG AGC GTG GGT GAT CCA  
ALA ARG HIS ASP ALA PRO ALA ARG HIS  
                  270                   275

15                   1400                   1410                   1420                   1430                   1440  
GCC GGA GAT CCA TCC GGA CCG GCG CGG CGA GTT CCA CGC GTG GTT CCA

20                   1450                   1460                   1470                   1480  
GAG CCA GCC GAG TTC CGG CGG CTG ACC GGT CAC TCC TTC TCC GTG CCG

                  1490                   1500                   1510                   1520                   1530  
CAG GTC GTC AAT ATC GCG TGT CCC GGA AAG GCG CCG CTG CGG CAT CCA

25                   1540                   1550                   1560                   1570                   1580  
CTT CTG CCG AGG TGC CAC CGG GCC GAG GCC AAG TAC AGC GGC GTG TGT

                  1590                   1600                   1610                   1620                   1630  
GCA GGG CGC CGG TGT CGA GGT CGT CGT CGA CGC GCC GGT GTC GAG GTC

30                   1640  
GTC GTC GAC-3'  
wherein A is deoxyadenyl residue; G is a deoxyquanyl residue; C  
is a deoxycytidyl residue; and T is a deoxythymidyl residue.

2309

87265660

0238323

26. The amino acid sequence encoded by nucleotides 541 to 1371 of the tylF gene defined in claim 25.


5

2310

87265660

# FIG. I


## The Tylosin Biosynthetic Pathway



87265660


## FIG.2

Restriction Site and Function Map of  
Plasmid pHJL280  
(26.94 kb)



## FIG.3


Restriction Site and Function Map of  
Plasmid pHJL284  
(26.94 kb)






## FIG.4

Restriction Site and Function Map of  
Plasmid pHJL309  
(28.24 kb)




## FIG.5

Restriction Site and Function Map of  
Plasmid pHJL311  
(30.84 kb)




## FIG.6

Restriction Site and Function Map of  
Plasmid pHJL315  
(38.83 kb)



**FIG.7**  
Chromosomal Organization of the Tylosin Biosynthetic Genes



87265660

2317